

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number
WO 00/78336 A1

- (51) International Patent Classification⁷: **A61K 38/16**, 38/20
- (21) International Application Number: PCT/US00/17103
- (22) International Filing Date: 21 June 2000 (21.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/334,512 21 June 1999 (21.06.1999) US
- (71) Applicants: **GENETICS INSTITUTE, INC.** [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US). **JOHNS HOPKINS UNIVERSITY** [US/US]; 615 North Wolfe Street, Baltimore, MD 21205 (US).
- (72) Inventors: **COLLINS, Mary**; 54 Rathburn Road, Natick, MA 01760 (US). **DONALDSON, Debra**; 108 Blakely, Medford, MA 02155 (US). **FITZ, Lori**; 13 Palmer Street, Arlington, MA 02174 (US). **NEBEN, Tamlyn**; 1115 Danforth Lane, Walnut Creek, CA 94598 (US). **WHITTERS, Matthew, J.**; 14 Brenton Wood Road, Hudson, MA 01749 (US). **WOOD, Clive**; 2 Hawthorne Place, #17R, Boston, MA 02114 (US). **WILLS-KARP, Marsha**; Johns Hopkins University, 615 North Wolfe Street, Baltimore, MD 21205 (US).
- (74) Agent: **HEHMAN, M., Kymne**; American Home products Corporation, Patent & Trademark Department - 2B, Attn: Kay E. Brady, One Campus Drive, Parsippany, NJ 07054 (US).
- (81) Designated States (*national*): AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TREATMENT OF FIBROSIS BY ANTAGONISM OF IL-13 AND IL-13 RECEPTOR CHAINS

(57) Abstract: Methods are provided for treating or inhibiting the formation of tissue fibrosis using IL-13 antagonists, including without limitation soluble forms of the IL-13 receptor.

WO 00/78336 A1

TREATMENT OF FIBROSIS
BY ANTAGONISM OF IL-13 AND IL-13 RECEPTOR CHAINS

5

This application is a continuation-in-part of application Ser. No. 09/301,808, filed April 28, 1999; which is a continuation-in-part of application Ser. No. 08/841,751, filed April 30, 1997; which is a divisional application of application Ser. No. 08/609,572, filed March 1, 1996 and issued as U.S. Patent No. 5,710,023 on January 20, 1998; all of which
10 are incorporated by reference herein.

Field of the Invention

The present invention relates to the treatment and inhibition of fibrosis by antagonism of the interaction of IL-13 with its receptor and receptor components.

15

Background of the Invention

A variety of regulatory molecules, known as cytokines, have been identified including interleukin-13 (IL-13). Various protein forms of IL-13 and DNA encoding various forms of IL-13 activity are described in McKenzie et al., Proc. Natl. Acad. Sci. USA 90:3735 (1993); Minty et al., Nature 362:248 (1993); and Aversa et al.,
20 WO94/04680. Thus, the term "IL-13" includes proteins having the sequence and/or biological activity described in these documents, whether produced by recombinant genetic engineering techniques; purified from cell sources producing the factor naturally or upon induction with other factors; or synthesized by chemical techniques; or a combination of
25 the foregoing.

IL-13 is a cytokine that has been implicated in production of several biological activities including: induction of IgG4 and IgE switching, including in human immature B cells (Punnonen et al., J. Immunol. 152:1094 (1994)); induction of germ line IgE heavy chain (ε) transcription and CD23 expression in normal human B cells (Punnonen et al.,
30 Proc. Natl. Acad. Sci. USA 90:3730 (1993)); and induction of B cell proliferation in the presence of CD40L or anti-CD40 mAb (Cocks et al., Int. Immunol. 5:657 (1993)). Although many activities of IL-13 are similar to those of IL-4, in contrast to IL-4, IL-13

does not have growth promoting effects on activated T cells or T cell clones (Zurawski et al., EMBO J. 12:2663 (1993)).

Like most cytokines, IL-13 exhibits certain biological activities by interacting with an IL-13 receptor ("IL-13R") on the surface of target cells. IL-13R and the IL-4 receptor
5 ("IL-4R") sharing a common component, which is required for receptor activation; however, IL-13 does not bind to cells transfected with the 130 kD IL-4R (Zurawski et al., *supra*). Thus, the IL-13R must contain at least one other ligand binding chain. Cytokine receptors are commonly composed of two or three chains. The cloning of one ligand binding chain for IL-13 has been recently reported (Hilton et al., Proc. Natl. Acad. Sci.
10 93:497-501).

It would be desirable to identify and clone the sequence for any other IL-13 binding chain of IL-13R so that IL-13R proteins can be produced for various reasons, including production of therapeutics and screening for inhibitors of IL-13 binding to the receptor and receptor signaling.

15

Summary of the Invention

In accordance with the present invention, polynucleotides encoding the IL-13 binding chains of the interleukin-13 receptor are disclosed, including without limitation those from the murine and human receptors. In certain embodiments, the invention
20 provides an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 256 to nucleotide 1404;
- (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 103 to nucleotide
25 1242;
- (c) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) or (b) as a result of degeneracy of the genetic code;
- (d) a nucleotide sequence capable of hybridizing under stringent conditions to the nucleotide specified in (a) or (b);
- 30 (e) a nucleotide sequence encoding a species homologue of the sequence specified in (a) or (b); and
- (f) an allelic variant of the nucleotide sequence specified in (a) or (b).

Preferably, the nucleotide sequence encodes a protein having a biological activity of the human IL-13 receptor. The nucleotide sequence may be operably linked to an expression control sequence. In preferred embodiments, the polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 256 to nucleotide 1404; the nucleotide
 5 sequence of SEQ ID NO:1 from nucleotide 319 to nucleotide 1257; the nucleotide sequence of SEQ ID NO:1 from nucleotide 1324 to nucleotide 1404; the nucleotide sequence of SEQ ID NO:3 from nucleotide 103 to nucleotide 1242; the nucleotide sequence of SEQ ID NO:3 from nucleotide 178 to nucleotide 1125; or the nucleotide sequence of SEQ ID NO:3 from nucleotide 1189 to nucleotide 1242.

10 The invention also provides isolated polynucleotides comprising a nucleotide sequence encoding a peptide or protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
 - (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
 - 15 (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
 - (d) the amino acid sequence of SEQ ID NO:4;
 - (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
 - (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
 - (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor
- 20 binding chain. Other preferred embodiments encode the amino acid sequence of SEQ ID NO:2 from amino acids 1 to 331 and the amino acid sequence of SEQ ID NO:2 from amino acids 26 to 331.

Host cells, preferably mammalian cells, transformed with the polynucleotides are also provided.

25 In other embodiments, the invention provides a process for producing a IL-13bc protein. The process comprises:

- (a) growing a culture of the host cell of the present invention in a suitable culture medium; and
- (b) purifying the human IL-13bc protein from the culture.

30 Proteins produced according to these methods are also provided.

The present invention also provides for an isolated IL-13bc protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- 5 (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain

Preferably the protein comprises the amino acid sequence of SEQ ID NO:2; the sequence
10 from amino acid 22 to 334 of SEQ ID NO:2; the sequence of SEQ ID NO:4; or the sequence from amino acid 26 to 341 of SEQ ID NO:4. In other preferred embodiments, the specified amino acid sequence is part of a fusion protein (with an additional amino acid sequence not derived from IL-13bc). Preferred fusion proteins comprise an antibody fragment, such as an Fc fragment. Particularly preferred embodiments comprise the amino
15 acid sequence of SEQ ID NO:2 from amino acids 1 to 331 and the amino acid sequence of SEQ ID NO:2 from amino acids 26 to 331.

Pharmaceutical compositions comprising a protein of the present invention and a pharmaceutically acceptable carrier are also provided.

The present invention further provides for compositions comprising an antibody
20 which specifically reacts with a protein of the present invention.

Methods of identifying an inhibitor of IL-13 binding to the IL-13bc or IL-13 receptor are also provided. These methods comprise:

- (a) combining an IL-13bc protein or a fragment thereof with IL-13 or a fragment thereof, said combination forming a first binding mixture;
- 25 (b) measuring the amount of binding between the protein and the IL-13 or fragment in the first binding mixture;
- (c) combining a compound with the protein and the IL-13 or fragment to form a second binding mixture;
- (d) measuring the amount of binding in the second binding mixture; and
- 30 (e) comparing the amount of binding in the first binding mixture with the amount of binding in the second binding mixture;

wherein the compound is capable of inhibiting IL-13 binding to the IL-13bc protein or IL-13 receptor when a decrease in the amount of binding of the second binding mixture occurs. Inhibitors of IL-13R identified by these methods and pharmaceutical compositions containing them are also provided.

- 5 Methods of inhibiting binding of IL-13 to the IL-13bc proteins or IL-13 receptor in a mammalian subject are also disclosed which comprise administering a therapeutically effective amount of a composition containing an IL-13bc protein, an IL-13bc or IL-13R inhibitor or an antibody to an IL-13bc protein.

10 Methods are also provided for potentiating IL-13 activity, which comprise combining a protein having IL-13 activity with a protein of the present invention and contacting such combination with a cell expressing at least one chain of IL-13R other than IL-13bc. Preferably, the contacting step is performed by administering a therapeutically effective amount of such combination to a mammalian subject.

15 Further methods are provided for treating an IL-13-related condition in a mammalian subject, said method comprising administering a therapeutically effective amount of a composition comprising an IL-13 antagonist and a pharmaceutically acceptable carrier. Other methods provide for a method of inhibiting the interaction of IL-13 with an IL-13bc protein in a mammalian subject comprising administering a therapeutically effective amount of a composition comprising an IL-13 antagonist and a pharmaceutically acceptable carrier. Preferably, the antagonist is selected from the group consisting of an IL-13bc protein, a soluble form of IL-13R α 1, an antibody to IL-13 or an IL-13-binding fragment thereof, an antibody to IL-13bc or an IL-13bc-binding fragment thereof, an antibody to IL-13R α 1 or an IL-13R α 1-binding fragment thereof, IL-13R-binding mutants of IL-4, a small molecule capable of inhibiting the interaction of IL-13 with IL-13bc and a small molecule capable of inhibiting the interaction of IL-13 with IL-13R α 1.

25 In yet other embodiments, the invention provides for a method of treating tissue fibrosis in a mammalian subject. The method comprises administering a therapeutically effective amount of a pharmaceutical composition comprising a protein and a pharmaceutically acceptable carrier, wherein the protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;

- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- 5 (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain.

The invention also provides for a method of inhibiting formation of tissue fibrosis in a mammalian subject. The method comprises administering a therapeutically effective
10 amount of a pharmaceutical composition comprising a protein and a pharmaceutically acceptable carrier, wherein the protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- 15 (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding
20 chain.

Other embodiments of the invention provide for a method of treating or inhibiting tissue fibrosis in a mammalian subject. The method comprises administering a therapeutically effective amount of a composition comprising (a) a molecule selected from the group consisting of an IL-13 antagonist and an IL-4 antagonist, and (b) a
25 pharmaceutically acceptable carrier.

In practicing such methods of treating or inhibiting fibrosis, preferably the tissue fibrosis affects a tissue selected from the group consisting of liver, skin epidermis, skin endodermis, muscle, tendon, cartilage, cardiac tissue, pancreatic tissue, lung tissue, uterine tissue, neural tissue, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary
30 tract and gut; most preferably, liver tissue (including tissue infected with schistosoma). In certain embodiments, the fibrosis results from the healing of a wound (including a surgical incision).

In practicing such methods of treating or inhibiting fibrosis using an antagonist, preferably such antagonist is selected from the group consisting of an IL-13bc protein, a soluble form of IL-13R α 1, an antibody to IL-13 or an IL-13-binding fragment thereof, an antibody to IL-13bc or an IL-13bc-binding fragment thereof, an antibody to IL-13R α 1 or
 5 an IL-13R α 1-binding fragment thereof, IL-13R-binding mutants of IL-4, a small molecule capable of inhibiting the interaction of IL-13 with IL-13bc and a small molecule capable of inhibiting the interaction of IL-13 with IL-13R α 1. In particularly preferred embodiments, the antagonist is an IL-13bc protein comprising an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- 15 (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain.

In other preferred methods of practicing such methods using an antagonist the antagonist is selected from the group consisting of a soluble form of IL-4R, an antibody
 20 to IL-4 or an IL-4-binding fragment thereof, an antibody to IL-4R or an IL-4R-binding fragment thereof, and a small molecule capable of inhibiting the interaction of IL-4 with IL-4R.

Brief Description of the Drawings

25 Fig. 1: The figure presents photographs of IL-13, IL-4, IL-11 and mock transfected COS cells after exposure to IL-13bc-Fc as described in Example 4 below.

Fig. 2: Characterization of the roles of IL-4 and IL-13 in schistosomiasis pathogenesis. C57BL/6 WT and IL-4-deficient (4KO) mice were infected with 25 cercariae of *Schistosoma mansoni* and then sacrificed at week 8 post-infection to evaluate the size
 30 of liver granulomas (panel A), tissue eosinophilia (panel B), and hepatic fibrosis (panel C). Separate groups of mice were treated with control-Fc or sIL-13R α 2-Fc as described in the Methods section. The data shown are measurements from individual mice and the lines

designate the means for each group. Statistical comparisons were made by Student's t-test (panels A and B) and by Analysis of Covariance (panel C). Significant comparisons and their p values are indicated in the figure. All data were reproduced in a second study.

Fig. 3: Liver collagen is reduced in sIL-13R α 2-Fc-treated /infected mice. Liver sections were prepared 8 weeks after challenge infection. Sections from control Fc-treated (panels A and B) and sIL-13R α 2-Fc-treated WT infected mice that contained nearly identical tissue egg burdens were stained with picosirius red (panels A and C) and illuminated using polarized light to highlight the areas rich in collagen (panels B and D). Birefringent areas indicate positive collagen staining and the areas shown are representative for each liver (magnification, X 40). Liver sections from sIL-13R α 2-Fc-treated mice showed only very slight granuloma and portal tract-associated collagen, in comparison with control animals.

Fig. 4: The Th1/Th2-type cytokine profile is unaffected by sIL-13R α 2-Fc treatment. C57BL/6 WT and IL-4-deficient (4KO) mice were infected with 25 cercariae of *Schistosoma mansoni* and separate groups of mice were treated with control-Fc or sIL-13R α 2-Fc as described in the Methods section. Mesenteric lymph node cells were isolated from individual mice and single cell suspensions were prepared (3×10^6 cells/well in 24 well plates) and stimulated with medium alone (squares), SEA at 20 ug/ml (circles), or with SEA and 50 ug/ml of anti-CD4 mAb (triangles). All cytokines were assayed in culture supernatants by ELISA 72 hrs post-stimulation as described in the Methods section. The symbols represent values for individual mice and the bars indicate the means within each group.

Fig. 5: Th2-type cytokine mRNA expression is reduced in the livers of infected IL-4-deficient mice but unaffected by IL-13 blockade. C57BL/6 WT and IL-4-deficient (4KO) mice were infected with 25 cercariae of *Schistosoma mansoni* and separate groups of mice were treated with control-Fc or sIL-13R α 2-Fc as described in the Methods section. All animals were sacrificed on wk 8 postinfection and liver specimens were prepared for RT-PCR analysis as described in the Methods section. The data shown are the individual values of 9 to 10 animals per group and the bar indicates the average within each group. The * symbol indicates that the data are significantly different from the WT control-Fc group as determined by Student's t-test ($p < .05$). The average values from five uninfected

WT (black circle) and five uninfected IL-4-deficient mice (open circle) are shown on the Y-axis for each cytokine. All data were reproduced in a second study.

Fig. 6: Collagen I and Collagen III mRNA expression is reduced in the livers of infected sIL-13R α 2-Fc treated mice, but unaffected by IL-4-deficiency. C57BL/6 WT and IL-4-deficient (4KO) mice were infected with 25 cercariae of *Schistosoma mansoni* and separate groups of mice were treated with control-Fc or sIL-13R α 2-Fc as described in the Methods section. All animals were sacrificed on wk 8 postinfection and liver specimens were prepared for RT-PCR analysis as described in the Methods section. The data shown are the individual values of 9 to 10 animals per group and the bar indicates the average within each group. The * symbol indicates that the data are significantly different from the WT and IL-4-deficient control-Fc groups as determined by Student's t-test ($p < .05$). The average values from five uninfected WT (black circle) and five uninfected IL-4-deficient mice (open circle) are shown on the Y-axis for each cytokine. These data were reproduced in a separate study.

Fig. 7: IL-13 induces type I collagen synthesis in murine 3T3 fibroblasts. Cells were stimulated with media (lane 1), rIL-4 at 1000 Units/ml (lane 2) or rIL-13 at 20 ng/ml (lanes 3 and 4, from R&D Systems and Genetics Institute, respectively) for 48 h. Total cell lysates were separated on 6% SDS-PAGE under reducing conditions, transferred to nitrocellulose membrane and probed with rabbit IgG anti-mouse type I collagen. The top doublet and bottom band (arrows) correspond to the purified rat collagen type I separated in lane 5 (panel A). The bottom figure (panel B) is the densitometric values (arbitrary pixel units).

Detailed Description of Preferred Embodiments

The inventors of the present application have for the first time identified and provided polynucleotides encoding the IL-13 binding chain of IL-13R (hereinafter "IL-13bc"), including without limitation polynucleotides encoding murine and human IL-13bc.

SEQ ID NO:1 provides the nucleotide sequence of a cDNA encoding the murine IL-13bc. SEQ ID NO:2 provides predicted the amino acid sequence of the receptor chain, including a putative signal sequence from amino acids 1-21. The mature murine IL-13bc is believed to have the sequence of amino acids 22-383 of SEQ ID NO:2. The mature murine receptor chain has at least three distinct domains: an extracellular domain

(comprising approximately amino acids 22-334 of SEQ ID NO:2), a transmembrane domain (comprising approximately amino acids 335-356 of SEQ ID NO:2) and an intracellular domain (comprising approximately amino acids 357-383 of SEQ ID NO:2).

SEQ ID NO:3 provides the nucleotide sequence of a cDNA encoding the human
5 IL-13bc. SEQ ID NO:4 provides predicted the amino acid sequence of the receptor chain, including a putative signal sequence from amino acids 1-25. The mature human IL-13bc is believed to have the sequence of amino acids 26-380 of SEQ ID NO:4. The mature human receptor chain has at least three distinct domains: an extracellular domain (comprising approximately amino acids 26-341 of SEQ ID NO:4), a transmembrane
10 domain (comprising approximately amino acids 342-362 of SEQ ID NO:4) and an intracellular domain (comprising approximately amino acids 363-380 of SEQ ID NO:4).

The first 81 amino acids of the human IL-13bc sequence are identical to the translated sequence of an expressed sequence tag (EST) identified as "yg99f10.r1 Homo sapiens cDNA clone 41648 5'" and assigned database accession number R52795.gb_est2.
15 There are no homologies or sequence motifs in this EST sequence which would lead those skilled in the art to identify the encoded protein as a cytokine receptor. A cDNA clone corresponding to this database entry is publicly-available from the I.M.A.G.E. Consortium. Subsequent to the priority date of the present application, such clone was ordered by applicants and sequenced. The sequence of such clone was determined to be the sequence
20 previously reported by applicants as SEQ ID NO:3 herein.

Soluble forms of IL-13bc protein can also be produced. Such soluble forms include without limitation proteins comprising amino acids 1-334 or 22-334 of SEQ ID NO:2 or amino acids 1-341 or 26-341 of SEQ ID NO:4. The soluble forms of the IL-13bc are further characterized by being soluble in aqueous solution, preferably at room
25 temperature. IL-13bc proteins comprising only the intracellular domain or a portion thereof may also be produced. Any forms of IL-13bc of less than full length are encompassed within the present invention and are referred to herein collectively with full length and mature forms as "IL-13bc" or "IL-13bc proteins." IL-13bc proteins of less than full length may be produced by expressing a corresponding fragment of the polynucleotide
30 encoding the full-length IL-13bc protein (SEQ ID NO:1 or SEQ ID NO:3). These corresponding polynucleotide fragments are also part of the present invention. Modified polynucleotides as described above may be made by standard molecular biology

techniques, including construction of appropriate desired deletion mutants, site-directed mutagenesis methods or by the polymerase chain reaction using appropriate oligonucleotide primers.

For the purposes of the present invention, a protein has "a biological activity of the IL-13 receptor binding chain" if it possess one or more of the following characteristics: (1) the ability to bind IL-13 or a fragment thereof (preferably a biologically active fragment thereof); and/or (2) the ability to interact with the second non-IL-13-binding chain of IL-13R to produce a signal characteristic of the binding of IL-13 to IL-13R. Preferably, the biological activity possessed by the protein is the ability to bind IL-13 or a fragment hereof, more preferably with a K_D of about 0.1 to about 100 nM. Methods for determining whether a particular protein or peptide has such activity include without limitation the methods described in the examples provided herein.

IL-13bc or active fragments thereof (IL-13bc proteins) may be fused to carrier molecules such as immunoglobulins. For example, soluble forms of the IL-13bc may be fused through "linker" sequences to the Fc portion of an immunoglobulin. Other fusions proteins, such as those with GST, Lex-A or MBP, may also be used.

The invention also encompasses allelic variants of the nucleotide sequences as set forth in SEQ ID NO:1 or SEQ ID NO:3, that is, naturally-occurring alternative forms of the isolated polynucleotide of SEQ ID NO:1 or SEQ ID NO:3 which also encode IL-13bc proteins, preferably those proteins having a biological activity of IL-13bc. Also included in the invention are isolated polynucleotides which hybridize to the nucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3 under highly stringent conditions (for example, 0.1xSSC at 65°C). Isolated polynucleotides which encode IL-13bc proteins but which differ from the nucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3 by virtue of the degeneracy of the genetic code are also encompassed by the present invention. Variations in the nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3 which are caused by point mutations or by induced modifications are also included in the invention.

The present invention also provides polynucleotides encoding homologues of the murine and human IL-13bc from other animal species, particularly other mammalian species. Species homologues can be identified and isolated by making probes or primers from the murine or human sequences disclosed herein and screening a library from an

appropriate species, such as for example libraries constructed from PBMCs, thymus or testis of the relevant species.

The isolated polynucleotides of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in
5 Kaufman et al., Nucleic Acids Res. 19, 4485-4490 (1991), in order to produce the IL-13bc protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" means enzymatically or chemically ligated to form a covalent bond
10 between the isolated polynucleotide of the invention and the expression control sequence, in such a way that the IL-13bc protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the IL-13bc protein. Any cell type capable of expressing functional IL-13bc protein may be used.
15 Suitable mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK, Rat2, BaF3, 32D, FDCP-1, PC12, M1x or C2C12 cells.

20 The IL-13bc protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are
25 well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. Soluble forms of the IL-13bc protein may also be produced in insect cells using appropriate isolated polynucleotides as described above.

Alternatively, the IL-13bc protein may be produced in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Suitable bacterial strains include
5 *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins.

Expression in bacteria may result in formation of inclusion bodies incorporating the recombinant protein. Thus, refolding of the recombinant protein may be required in order to produce active or more active material. Several methods for obtaining correctly
10 folded heterologous proteins from bacterial inclusion bodies are known in the art. These methods generally involve solubilizing the protein from the inclusion bodies, then denaturing the protein completely using a chaotropic agent. When cysteine residues are present in the primary amino acid sequence of the protein, it is often necessary to accomplish the refolding in an environment which allows correct formation of disulfide
15 bonds (a redox system). General methods of refolding are disclosed in Kohno, Meth. Enzym., 185:187-195 (1990). EP 0433225 and copending application USSN 08/163,877 describe other appropriate methods.

The IL-13bc protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or
20 sheep which are characterized by somatic or germ cells containing a polynucleotide sequence encoding the IL-13bc protein.

The IL-13bc protein of the invention may be prepared by growing a culture transformed host cells under culture conditions necessary to express the desired protein. The resulting expressed protein may then be purified from the culture medium or cell
25 extracts. Soluble forms of the IL-13bc protein of the invention can be purified from conditioned media. Membrane-bound forms of IL-13bc protein of the invention can be purified by preparing a total membrane fraction from the expressing cell and extracting the membranes with a non-ionic detergent such as Triton X-100.

The IL-13bc protein can be purified using methods known to those skilled in the
30 art. For example, the IL-13bc protein of the invention can be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be

applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) or polyethyleimine (PEI) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-Sepharose® columns). The purification of the IL-13bc protein from culture supernatant may also include one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the IL-13bc protein. Affinity columns including IL-13 or fragments thereof or including antibodies to the IL-13bc protein can also be used in purification in accordance with known methods. Some or all of the foregoing purification steps, in various combinations or with other known methods, can also be employed to provide a substantially purified isolated recombinant protein. Preferably, the isolated IL-13bc protein is purified so that it is substantially free of other mammalian proteins.

IL-13bc proteins of the invention may also be used to screen for agents which are capable of binding to IL-13bc or IL-13R or which interfere with the binding of IL-13 to the IL-13 or IL-13bc (either the extracellular or intracellular domains) and thus may act as inhibitors of normal binding and cytokine action (“IL-13R inhibitors”). Binding assays using a desired binding protein, immobilized or not, are well known in the art and may be used for this purpose using the IL-13bc protein of the invention. Purified cell based or protein based (cell free) screening assays may be used to identify such agents. For example, IL-13bc protein may be immobilized in purified form on a carrier and binding to purified IL-13bc protein may be measured in the presence and in the absence of potential inhibiting agents. A suitable binding assay may alternatively employ a soluble form of IL-13bc of the invention. Another example of a system in which inhibitors may be screened is described in Example 2 below.

In such a screening assay, a first binding mixture is formed by combining IL-13 or a fragment thereof and IL-13bc protein, and the amount of binding in the first binding mixture (B_0) is measured. A second binding mixture is also formed by combining IL-13 or a fragment thereof, IL-13bc protein, and the compound or agent to be screened, and the amount of binding in the second binding mixture (B) is measured. The amounts of binding in the first and second binding mixtures are compared, for example, by performing a calculation of the ratio B/B_0 . A compound or agent is considered to be capable of inhibiting binding if a decrease in binding in the second binding mixture as compared to the first binding mixture is observed. Optionally, the second chain of IL-13R can be added to one or both of the binding mixtures. The formulation and optimization of binding mixtures is within the level of skill in the art, such binding mixtures may also contain buffers and salts necessary to enhance or to optimize binding, and additional control assays may be included in the screening assay of the invention.

Compounds found to reduce the binding activity of IL-13bc protein to IL-13 or its fragment to any degree, preferably by at least about 10%, more preferably greater than about 50% or more, may thus be identified and then secondarily screened in other binding assays and in vivo assays. By these means compounds having inhibitory activity for IL-13bc binding which may be suitable as therapeutic agents may be identified.

IL-13bc proteins, and polynucleotides encoding them, may also be used as diagnostic agents for detecting the expression or presence of IL-13bc, IL-13R, IL-13 or cells expressing IL-13bc, IL-13R or IL-13. The proteins or polynucleotides may be employed for such purpose in standard procedures for diagnostics assays using these types of materials. Suitable methods are well known to those skilled in the art.

As used herein "IL-13R" refers to IL-13bc and/or a second IL-13 receptor chain known as "IL-13R α 1" or "NR4" (see: murine receptor chain, Hilton et al., Proc. Natl. Acad. Sci. USA 1996, 93:497-501; human receptor chain, Aman et al., J. Biol. Chem. 1996, 271:29265-70, and Gauchat et al., Eur. J. Immunol. 1997, 27:971-8).

IL-13bc acts as a mediator of the known biological activities of IL-13. As a result, IL-13bc protein (particularly, soluble IL-13bc proteins), IL-13R inhibitors (i.e., antagonists of interaction of IL-13 with IL-13R (such as, for example, antibodies to IL-13R (including particularly to IL-13bc or to IL-13R α 1) and fragments thereof, antibodies to IL-13 and fragments thereof, soluble IL-13R α 1 proteins, and small molecule and other inhibitors of

the interaction of IL-13 with IL-13R (including with IL-13bc and/or with IL-13R α 1)) may be useful in treatment or modulation of various medical conditions in which IL-13 is implicated or which are effected by the activity (or lack thereof) of IL-13 (collectively "IL-13-related conditions"). Mutated forms of IL-4 which bind to IL-13R can also be used as

5 IL-13 antagonists (see, for example, those disclosed in Shanafelt et al., Proc. Natl. Acad. Sci. USA 1998, 95:9454-8; Aversa et al., J. Exp. Med. 1993, 178:2213-8; and Grunewald et al., J. Immunol. 1998, 160:4004-9).

IL-13-related conditions include without limitation Ig-mediated conditions and diseases, particularly IgE-mediated conditions (including without limitation atopy, allergic

10 conditions, asthma, immune complex diseases (such as, for example, lupus, nephrotic syndrome, nephritis, glomerulonephritis, thyroiditis and Grave's disease)); inflammatory conditions of the lungs; immune deficiencies, specifically deficiencies in hematopoietic progenitor cells, or disorders relating thereto; cancer and other disease. Such pathological states may result from disease, exposure to radiation or drugs, and include, for example,

15 leukopenia, bacterial and viral infections, anemia, B cell or T cell deficiencies such as immune cell or hematopoietic cell deficiency following a bone marrow transplantation. Since IL-13 inhibits macrophage activation, IL-13bc proteins may also be useful to enhance macrophage activation (i.e., in vaccination, treatment of mycobacterial or intracellular organisms, or parasitic infections).

20 IL-13bc proteins may also be used to potentiate the effects of IL-13 in vitro and in vivo. For example, an IL-13bc protein can be combined with a protein having IL-13 activity (preferably IL-13) and the resulting combination can be contacted with a cell expressing at least one chain of IL-13R other than IL-13bc (preferably all chains of IL-13R other than IL-13bc, such as IL-13R α 1). Preferably, the contacting step is performed by

25 administering a therapeutically effective amount of such combination to a mammalian subject in vivo. The pre-established association of the IL-13 protein with the IL-13bc protein will aid in formation of the complete IL-13/IL-13R complex necessary for proper signaling. See for example the methods described by Economides et al., Science 270:1351 (1995).

30 IL-13bc protein and IL-13R inhibitors, purified from cells or recombinantly produced, may be used as a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may contain, in addition to IL-

13bc or inhibitor and carrier, various diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of
5 administration.

The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, interleukins (such as, IL-1, IL-2, IL-3, IL-4 . . . IL-24, IL-25), G-CSF, stem cell factor, and erythropoietin. The pharmaceutical composition may also include anti-cytokine antibodies. The
10 pharmaceutical composition may further contain other anti-inflammatory agents. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with isolated IL-13bc protein or IL-13bc inhibitor, or to minimize side effects caused by the isolated IL-13bc or IL-13bc inhibitor. Conversely, isolated IL-13bc or IL-13bc inhibitor may be included in formulations of the particular
15 cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

The pharmaceutical composition of the invention may be in the form of a liposome in which isolated IL-13bc protein or IL-13bc inhibitor is combined, in addition to other
20 pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of
25 skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient
30 to show a meaningful patient benefit, e.g., amelioration of symptoms of, healing of, or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to

a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of isolated IL-13bc protein or IL-13bc inhibitor is administered to a mammal. Isolated IL-13bc protein or IL-13bc inhibitor may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, IL-13bc protein or IL-13bc inhibitor may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering IL-13bc protein or IL-13bc inhibitor in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of IL-13bc protein or IL-13bc inhibitor used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of IL-13bc protein or IL-13bc inhibitor is administered orally, IL-13bc protein or IL-13bc inhibitor will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% IL-13bc protein or IL-13bc inhibitor, and preferably from about 25 to 90% IL-13bc protein or IL-13bc inhibitor. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to

90% by weight of IL-13bc protein or IL-13bc inhibitor, and preferably from about 1 to 50% IL-13bc protein or IL-13bc inhibitor.

When a therapeutically effective amount of IL-13bc protein or IL-13bc inhibitor is administered by intravenous, cutaneous or subcutaneous injection, IL-13bc protein or
5 IL-13bc inhibitor will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to IL-13bc protein or IL-13bc inhibitor an isotonic vehicle such as
10 Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additive known to those of skill in the art.

The amount of IL-13bc protein or IL-13bc inhibitor in the pharmaceutical
15 composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of IL-13bc protein or IL-13bc inhibitor with which to treat each individual patient. Initially, the attending physician will administer low doses of IL-13bc protein or IL-13bc inhibitor and observe
20 the patient's response. Larger doses of IL-13bc protein or IL-13bc inhibitor may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1 μ g to about 100 mg (preferably about 20 μ g to about 500 μ g) of IL-13bc
25 protein or IL-13bc inhibitor per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the IL-13bc protein or IL-13bc
30 inhibitor will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

IL-13bc proteins of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the IL-13bc protein and which may inhibit binding of IL-13 or fragments thereof to the receptor. Such antibodies may be obtained using the entire IL-13bc as an immunogen, or by using
5 fragments of IL-13bc, such as the soluble mature IL-13bc. Smaller fragments of the IL-13bc may also be used to immunize animals. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Additional peptide immunogens may be generated by replacing tyrosine residues with sulfated tyrosine residues. Methods for synthesizing
10 such peptides are known in the art, for example, as in R.P. Merrifield, J.Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, et al., FEBS Lett. 211, 10 (1987).

Neutralizing or non-neutralizing antibodies (preferably monoclonal antibodies) binding to IL-13bc protein may also be useful therapeutics for certain tumors and also in the treatment of conditions described above. These neutralizing monoclonal antibodies
15 may be capable of blocking IL-13 binding to the IL-13bc.

Example 1

Isolation of IL-13bc cDNAs

Isolation of the murine IL-13 receptor chain.

20 5 ug of polyA+ RNA was prepared from the thymuses of 6-8 week old C3H/HeJ mice. Double stranded, hemimethylated cDNA was prepared using Stratagene's cDNA synthesis kit according to manufacturers instructions. Briefly, the first strand was primed with an oligodT-Xho primer, and after second strand synthesis, EcoRI adapters were added, and the cDNA was digested with XhoI, and purified. The cDNA was ligated to the
25 XhoI-EcoRI sites of the Zap Express (Stratagene) lambda vector, and packaged using Gigapak II Gold packaging extracts (Stratagene) according to the manufacturers instructions. A library of 1.5×10^6 resulting recombinant phage was amplified following manufacturer's instructions. This library was screened with a degenerate 17mer oligonucleotide probe of the sequence KSRCTCCABK CRCTCCA (SEQ ID NO:5) (K
30 = G+T; S= C+G; R=A+G; B=C+G+T) using standard TMAC hybridization conditions as described (Current Protocols in Molecular Biology, Ausubel, et al., editors., John Wiley and Sons, 1995, section 6.4.3). Clone A25 was identified because it hybridized to the

17mer probe, but not to probes derived from known hematopoietin receptors. This clone was isolated in plasmid form from the ZapExpress vector as per manufacturers instruction, and the DNA sequence was determined. The DNA sequence encoded a novel member of the hematopoietin receptor family.

- 5 Clone A25 containing the polynucleotide having the sequence of SEQ ID NO:1 was deposited with ATCC as pA25pBKCMV at accession number 69997 on February 22, 1996.

Isolation of the human IL-13 receptor chain.

- 10 A partial fragment of the human homolog of the murine receptor was isolated by PCR using oligonucleotides derived from the murine sequence. cDNA was prepared from human testis polyA+ RNA that was obtained from Clontech. A DNA fragment of 274 base pairs was amplified from this cDNA by PCR with the following oligonucleotides: ATAGTTAAACCATTGCCACC (SEQ ID NO:6) and
- 15 CTCCATTCGCTCCAAATTCC (SEQ ID NO:7) using AmpliTaq polymerase (Promega) in 1X Taq buffer containing 1.5 mM MgCl₂ for 30 cycles of incubation (94°C x 1 minute, 42°C for 1 minute, and 72°C for 1 minute). The DNA sequence of this fragment was determined, and two oligonucleotides were prepared from an internal portion of this fragment with the following sequence: AGTCTATCTTACTTTTACTCG (SEQ ID NO:8)
- 20 and CATCTGAGCAATAAATATTAC (SEQ ID NO:9). These oligonucleotides were used as probes to screen a human testis cDNA library purchased from CLONTECH (cat #HL1161). Filters were hybridized at 52°C using standard 5XSSC hybridization conditions and washed in 2X SSC at 52°C. Twenty two clones were isolated that hybridized to both oligonucleotides in a screen of 400,000 clones. DNA sequence was
- 25 determined from four of the cDNA clones, and all encoded the same novel hematopoietin receptor. The predicted DNA sequence of the full length human receptor chain is shown as SEQ ID NO:3.

The human clone was deposited with ATCC as phA25#11pDR2 at accession number 69998 on February 22, 1996.

30

Example 2

Expression of Soluble IL-13bc Protein and

Assay of ActivityProduction and purification of soluble IL-13bc-Ig.

DNA encoding amino acids 1-331 of the extracellular domain of murine IL-13bc
 5 was fused to a spacer sequence encoding gly-ser-gly by PCR and ligated in frame with
 sequences encoding the hinge CH2 CH3 regions of human IgG1 of the COS-1 expression
 vector pED.Fc. IL-13bc-Ig was produced from DEAE-dextran transfected COS-1 cells
 and purified via protein A sepharose chromatography (Pharmacia).

10 B9 proliferation assay

Stimulation of proliferation of B9 cells (Aarden et al. Eur. J. Immunol. 1987.
 17:1411-1416) in response to IL-13 or IL-4 was measured by 3H-thymidine incorporation
 into DNA. Cells (5 x 10³/well) were seeded into 96 well plates with media containing
 growth factors at varying concentrations in the presence or absence of IL-13bc-Ig at
 15 1ug/ml. After incubation for 3 days 1uCi/well of 3H-thymidine was added and the cells
 incubated for an additional 4 hrs. Incorporated radioactivity was determined using a LKB
 1205 Plate reader.

The B9 cell line proliferated in response to IL-13, IL-4 or IL-6. Only responses to
 IL-13 were inhibited by the soluble IL-13bc-Ig, indicating that this receptor binds IL-13
 20 specifically, but not IL-4 or IL-6. The tables show cpm. Two separate experiments are
 shown.

Table I

25	cytokine dilution	IL-13 (3ng/ml)	IL-13 plus A25-Fc (1ug/ml)	IL-4 (20 ng/ml)	IL-4 plus A25-Fc (1 ug/ml)	Cos IL-6 (1/10,000)
	1	37734	1943	6443	6945	37887
	1/3	30398	1571	2680	2442	36500
	1/10	16101	1461	1767	1771	33335
	1/30	2148	1567	1619	1783	27271
30	1/100	1574	1419	1522	1576	18831
	1/300	1512	1531	1373	1577	7768

1/1000	1316	1392	1190	1474	2760
1/3000	1834	1994	1482	1819	1672

5 Table II

	cytokine dilution	IL-13 (3ng/ml)	IL-13 plus A25-Fc (5ug/ml)	IL-4 (20 ng/ml)	IL-4 plus A25-Fc (5ug/ml)	Cos IL-6 (1/10,000)	Cos IL-6 plus A25-Fc (5ug/ml)
	1	6413	295	1216	1158	6969	7703
	1/3	5432	281	518	656	7827	8804
10	1/10	2051	281	489	520	8345	10027
	1/30	506	319	279	476	8680	9114
	1/100	430	372	288	423	7426	10364
	1/300	330	287	323	420	5531	6254
	1/1000	326	389	348	nt	2524	nt
15	no cytokine	339	279	404	394	326	279

Example 3Direct Binding of Soluble IL-13bc to IL-13 Measured by Surface Plasmon Resonance
(Biacore Analysis).

5 A Biacore biosensor was used to measure directly the specific binding of IL-13 to purified IL-13bc-Ig (Pharmacia, Johnsson et al., 1991). Approximately 10,000 to 17,000 resonance units (RU) of purified IL-13bc-Ig, human IgG1 or irrelevant receptor were each covalently immobilized to different flow cells on the sensor chip as recommended by the manufacturer. (RU's are a reflection of the mass of protein bound to the sensor chip

10 surface.) Purified IL-13 was injected across the flow cells at 5 ul/min for 10 mins in the presence or absence of excess purified IL-13bc-Ig. Binding was quantified as the difference in RU before and after sample injection. Specific IL-13 binding of 481.9 RU was observed only for immobilized IL-13bc-Ig whereas coinjection of IL-13 plus IL-13bc-Ig resulted in no binding to the immobilized IL-13bc-Ig (4 RU). No IL-13 binding was

15 observed for either immobilized IgG or IL-11R-Ig (5.4 and 3.7 RU respectively).

20

Sample	IL-13bc-Ig (10,383 RU)	IgG control (13,399 RU)	IL-11R-Ig (17,182 RU)
100 ng/ml human IL-13	481.9 RU bound	5.4 RU bound	3.7 RU bound
100 ng/ml human IL-13 + soluble IL-13bc-Ig	4.0 RU bound	not tested	not tested

25

Example 4

Binding of IL-13 Expressed in COS Cells to

Labeled IL-13BC-Ig Fusion Protein:

COS in situ Detection of IL-13 with IL-13bc-Fc

5

Expression vectors for IL-13, IL-4, IL-11 or empty vector were transfected into COS-1 cells in duplicated plates via the DEAE-dextran method. Two days after transfection cells were washed twice in phosphate buffered saline (PBS) and fixed in the culture dish for 10' at 4° C with methanol. Following fixation cells were washed twice
10 with PBS then rinsed once with binding buffer (PBS, 1% (w/v) bovine serum albumin, .1% (w/v) sodium azide) and incubated for two hours at 4° C in binding buffer with IL-13bc-Fc at 1.0ug/ml or with relevant anti-cytokine antisera. Cells were washed twice with PBS and incubated at 4o C with shaking in alkaline phosphatase labeled Rabbit F(ab)2' anti-human IgG diluted 1:500 in binding buffer (for Fc fusion detection) or Rabbit F(ab)2'
15 anti-rat IgG (for anti-cytokine detection) . Cells were again washed twice in PBS. Alkaline phosphatase activity was visualized using nitro blue tetrazolium and 5-bromo-4-chloro-3-indolyl-phosphate.

Specific binding was visualized under the microscope. Only cells transfected with IL-13 showed specific binding to IL13bc-Ig. (See photograph of transfected cells, Figure
20 1).

Example 5

Other Systems for Determination Biological Activity of IL-13bc Protein

Other systems can be used to determine whether a specific IL-13bc protein exhibits
25 a "biological activity" of IL-13bc as defined herein. The following are examples of such systems.

Assays for IL-13 Binding

The ability of a IL-13bc protein to bind IL-13 or a fragment thereof can be
30 determine by any suitable assays which can detect such binding. Some suitable examples follow.

Binding of IL-13 to the extracellular region of the IL-13bc protein will specifically cause a rapid induction of phosphotyrosine on the receptor protein. Assays for ligand binding activity as measured by induction of phosphorylation are described below.

Alternatively, a IL-13bc protein (such as, for example, a soluble form of the
5 extracellular domain) is produced and used to detect IL-13 binding. For example, a DNA construct is prepared in which the extracellular domain (truncated prior, preferably immediately prior, to the predicted transmembrane domain) is ligated in frame to a cDNA encoding the hinge C_H2 and C_H3 domains of a human immunoglobulin (Ig) γ 1. This construct is generated in an appropriate expression vector for COS cells, such as pED Δ C
10 or pMT2. The plasmid is transiently transfected into COS cells. The secreted IL-13bc-Ig fusion protein is collected in the conditioned medium and purified by protein A chromatography.

The purified IL-13bc-Ig fusion protein is used to demonstrate IL-13 binding in a number of applications. IL-13 can be coated onto the surface of an enzyme-linked
15 immunosorbent assay (ELISA) plate, and then additional binding sites blocked with bovine serum albumin or casein using standard ELISA buffers. The IL-13bc-Ig fusion protein is then bound to the solid-phase IL-13, and binding is detected with a secondary goat anti-human Ig conjugated to horseradish peroxidase. The activity of specifically bound enzyme can be measured with a colorimetric substrate, such as tetramethyl benzidine and
20 absorbance readings.

IL-13 may also be expressed on the surface of cells, for example by providing a transmembrane domain or glucosyl phosphatidyl inositol (GPI) linkage. Cells expressing the membrane bound IL-13 can be identified using the IL-13bc-Ig fusion protein. The soluble IL-13bc-Ig fusion is bound to the surface of these cells and detected with goat anti-
25 human Ig conjugated to a fluorochrome, such as fluorescein isothiocyanate and flow cytometry.

Interaction Trap

A yeast genetic selection method, the "interaction trap" [Gyuris et al, Cell
30 75:791-803, 1993], can be used to determine whether a IL-13bc protein has a biological activity of IL-13bc as defined herein. In this system, the expression of reporter genes from both LexAop-Leu2 and LexAop-LacZ relies on the interaction between the bait protein,

for example in this case a species which interacts with human IL-13bc, and the prey, for example in this case the human IL-13bc protein. Thus, one can measure the strength of the interaction by the level of Leu2 or LacZ expression. The most simple method is to measure the activity of the LacZ encoded protein, β -galactosidase. This activity can be judged by the degree of blueness on the X-Gal containing medium or filter. For the quantitative measurement of β -galactosidase activity, standard assays can be found in "Methods in Yeast Genetics" Cold Spring Harbor, New York, 1990 (by Rose, M.D., Winston, F., and Hieter, P.).

In such methods, if one wishes to determine whether the IL-13bc protein interacts with a particular species (such as, for example, a cytosolic protein which binds to the intracellular domain of the IL-13bc *in vivo*), that species can be used as the "bait" in the interaction trap with the IL-13bc protein to be tested serving as the "prey", or *vice versa*.

Example 6

Inhibition of Fibrosis Using Soluble IL-13R

The development of fibrous tissue is part of the normal process of healing after injury. Nevertheless, in some circumstances there is a destructive accumulation of excess collagen that interferes with the normal function of the affected tissue. Indeed collagen synthesis and tissue scarring are the major pathological manifestations of a number of chronic and debilitating illnesses, including several autoimmune, allergic, and infectious diseases¹⁻⁷. While there is a great deal of mechanistic information regarding the process of scar tissue formation^{8,9}, there are still large gaps in our understanding of the role of inflammatory cells and cytokines in initiating the fibrotic process.

As used herein "fibrosis" includes any condition which involves the formation of fibrous tissue (whether such formation is desirable or undesirable). Such conditions include, without limitation, fibrositis, formation of fibromas (fibromatosis), fibrogenesis (including pulmonary fibrogenesis), fibroelastosis (including endocardial fibroelastosis), formation of fibromyomas, fibrous ankylosis, formation of fibroids, formation of fibroadenomas, formation of fibromyxomas, and fibrocystitis (including cystic fibrosis).

The IL-13 receptor complex is composed of at least three distinct components, including the IL-4 receptor, the low-affinity binding IL-13Ra1 chain, and the high affinity

binding chain, IL-13R α 2^{35,42-44}. Recently, a soluble IL-13Ra2-Fc fusion protein was prepared and has been used successfully to neutralize IL-13 both *in vitro*³⁵ and *in vivo*^{30,39-41}. Since the fusion protein binds IL-13 with high affinity, but fails to neutralize IL-4, the protein provided an excellent tool to determine the specific roles of IL-13. In the present study, we used the IL-13 antagonist in wild type and IL-4-deficient mice in order to dissect the contributions of IL-13 and IL-4 to the development of hepatic fibrosis in murine schistosomiasis. In these studies, granuloma formation was examined in detail, focusing on eosinophil and mast cell recruitment and, more importantly, the development of egg-induced fibrosis was quantified using biochemical, histological, and molecular techniques. We also examined the contributions of IL-4 and IL-13 to the regulation of Th1/Th2-type cytokine responses both *in vitro*, in mesenteric lymph node cultures and, *in vivo*, in the granulomatous livers. While the results from this study show that IL-13 and IL-4 exhibit some redundant activities in schistosomiasis pathogenesis, distinct functions for both cytokines were also clearly elucidated. Probably the most important and novel finding was the observation that IL-13, not IL-4, was the major Th2-type cytokine driving type I and type III collagen mRNA production and hepatic fibrosis in infected mice. Thus, our findings establish that an IL-13 inhibitor/antagonist, such as sIL-13R α 2-Fc, can be of therapeutic benefit in preventing fibrosis, such as, for example, that associated with chronic infectious disease.

20

RESULTS

Comparative effect of IL-4, IL-13 or double IL-4/IL-13 deficiencies in schistosomiasis pathogenesis: sIL-13R α 2-Fc treatment significantly reduces hepatic fibrosis in S. mansoni-infected mice

To compare the regulatory roles of IL-4 and IL-13 in the pathogenesis of schistosomiasis, we infected C57BL/6 WT and IL-4-deficient mice percutaneously with 25 *S. mansoni* cercariae. Separate groups of animals were treated with either sIL-13R α 2-Fc or with control-Fc, as described in the Materials and Methods. The treatments began on week 5, at the start of egg laying, and all animals were sacrificed 8 wk postinfection and examined for several parasitologic and immunologic parameters. As shown in Table III,

30

{TABLE III INSERTED HERE}

all four groups of mice harbored similar worm burdens, and tissue eggs produced per worm pair did not vary among the groups. At 8 wk postinfection, the time of the peak tissue response⁴⁵, WT mice showed no significant change in granuloma size as a result of IL-13 blockade (Fig. 2A). Interestingly, control-Fc-treated IL-4-deficient mice also
5 failed to show a reduced granulomatous response, and in fact, granulomas were significantly larger in these mice. In striking contrast to these observations, the IL-4-deficient mice displayed a markedly reduced granulomatous response when IL-13 was inhibited (Fig. 2A, far right). Indeed, the double IL-4-deficient/sIL-13R α 2-Fc-treated mice displayed on average a 40 to 50% reduction in granuloma volume when compared
10 with either control or sIL-13R α 2-Fc-treated WT animals, and more than a 75% reduction when compared with control-Fc-treated IL-4-deficient mice.

The cellular composition of the lesions was also evaluated in Giemsa-stained liver sections and as shown in Table III, IL-4-deficient mice displayed a marked reduction in granuloma-associated mast cells. In contrast, there was no change in mast cell numbers by
15 IL-13 inhibition alone, and IL-13 blockade had no additional effect on the already highly reduced numbers of mast cells in IL-4-deficient mice. Somewhat similar, yet distinct findings were observed when granuloma-associated eosinophils were evaluated (Fig. 2B). Here, the numbers of eosinophils were increased from 46 to 64% in WT mice by IL-13 blockade and significantly decreased (28%) as consequence of IL-4 deficiency. Despite
20 the apparent contrasting roles for IL-13 and IL-4 in the tissue eosinophilia, an even more striking combined inhibitory effect was observed when the IL-4-deficient mice were treated with the IL-13 inhibitor. In these mice, the average number of granuloma eosinophils was below 10%. Finally, there was no change in the degree of parenchymal or egg-associated liver necrosis in the WT versus IL-4-deficient animals, while both sIL-
25 13R α 2-Fc-treated WT and IL-4-deficient groups showed marked reductions in overall parenchymal necrosis.

Perhaps most importantly, the sIL-13R α 2-Fc treatment alone significantly reduced the collagen content of liver granulomas in WT mice, as assessed in tissue sections stained with picrosirius red (Table III and Fig. 3). In contrast, infected IL-4-deficient mice showed
30 no detectable change in granuloma collagen deposition by microscopic analysis. Interestingly, there appeared to be no combined or synergistic role for IL-13 and IL-4 in this parameter since there was no significant difference between sIL-13R α 2-Fc-treated-

WT and -IL-4 deficient mice (Table III). Fig. 3 shows that while similar worm numbers, tissue egg burdens, and granuloma sizes were found in control and sIL-13R α 2-Fc treated WT mice, IL-13 blockade had a substantial inhibitory effect on collagen deposition within the liver. Finally, the extent of hepatic fibrosis was also measured by the assessment of hydroxyproline levels (Fig. 2C), which is more quantitative than the histological techniques described above. The soluble IL-13 antagonist alone markedly decreased liver hydroxyproline levels, while the IL-4-deficiency resulted in a less significant reduction. The dual IL-4/IL-13 deficiency failed to reduce hydroxyproline to levels below that already observed in the sIL-13R α 2-Fc treated WT mice (Fig. 2C), although there was a slight trend in a second study (not significant). Together, these data demonstrate that IL-13 is the dominant Th2-associated cytokine responsible for the development of hepatic fibrosis in murine schistosomiasis.

Th2-type cytokine production is reduced in IL-4-deficient mice but unaffected by IL-13 inhibition.

While it is well-known that IL-4 is the primary cytokine driving CD4⁺ Th2 cell development^{21,22}, the role of IL-13 in the generation and maintenance of Th2-type responses has been controversial and may be influenced by both host genetics and the infectious disease model under study^{30,34,38}. Therefore, to determine whether the sIL-13R α 2-Fc-induced changes in liver pathology were generated by alterations in the Th1/Th2 cytokine balance, we isolated mesenteric lymph nodes and spleens from infected mice, prepared single cell suspensions, and restimulated the cultures *in vitro* with parasite antigens. Additional cell cultures were exposed to parasite antigens in the presence of anti-CD4 mAbs to determine whether cytokine production was dependent upon a CD4⁺ T cell response. Culture supernatants were analyzed by ELISA for IL-4, IL-13, IL-5, IL-10, and IFN- γ . As might be predicted¹⁵, mesenteric (Fig. 5) and splenic cultures (data not shown) prepared from WT mice displayed a highly polarized Th2-type cytokine response. They produced high levels of IL-4, IL-5, IL-10, and IL-13 in response to SEA stimulation and little or no IFN- γ . IL-4-deficient mice in contrast showed a more mixed Th1/Th2-type profile. Indeed, a significant SEA-specific IFN- γ response was detected in IL-4-deficient mice, which is consistent with previous studies^{23,24}. IL-13, IL-10, and to a lesser extent IL-5, were also detected in these animals, although the levels of these cytokines were

markedly decreased when compared with WT mice. Importantly, the maintenance of the low but significant IL-4-independent IL-13 response likely explains the marked granulomatous response that is maintained in the absence IL-4 (Fig. 2). Surprisingly, despite its marked inhibitory effect on hepatic fibrosis, sIL-13R α 2-Fc had no significant effect on Th1 or Th2-type cytokine responses in either WT or IL-4-deficient mice. It should also be noted that in all cases, cytokine production was highly dependent on a CD4⁺ T cell response, since little or no cytokine expression was detected in any of the anti-CD4 mAb-treated SEA-stimulated cultures.

10 ***Changes in Th1/Th2-type cytokine mRNA expression in the granulomatous livers of IL-4-deficient and sIL-13R α 2-Fc-treated mice.***

To determine whether a similar pattern of cytokine expression was observed *in vivo* at the site of granuloma formation, we isolated liver mRNA from the various groups of mice at 8 wk postinfection and performed quantitative RT-PCR. As shown in Figure 5, infected WT mice displayed a strong Th2-type cytokine mRNA profile, showing marked increases in IL-4, IL-13, IL-5, and IL-10 mRNA. The WT mice also showed modest increases in the expression of IFN- γ mRNA, which was consistent with previous observations¹⁹. In contrast to these findings, IL-13 and IL-5 mRNA levels were much lower in IL-4-deficient mice, while IL-10 and TNF- α mRNA significantly increased and IFN- γ mRNA expression did not change. Again, similar to the *in vitro* results obtained from mesenteric lymph node and splenocyte cultures, IL-13 blockade had no significant effect on the pattern of cytokine mRNA expression in either WT or IL-4-deficient mice. There was however, a modest increase in IL-10 mRNA levels in IL-4-deficient mice treated with the sIL-13R α 2-Fc, although this is unlikely to explain the decreases in fibrosis, since highly divergent levels of IL-10 were detected in sIL-13R α 2-Fc-treated WT versus IL-4-deficient mice, yet a similar decrease in fibrosis was observed. TGF- β 1 and TGF- β 2 mRNA expression was also examined in the granulomatous tissues, however no significant differences were observed in either infected IL-4-deficient mice or in animals treated with sIL-13R α 2-Fc (data not shown).

30

Collagen I and collagen III mRNA levels are reduced in the livers of sIL-13R α 2-Fc-treated mice but unaffected by IL-4-deficiency.

The *in vitro* and *in vivo* cytokine studies described above suggested that the anti-fibrotic effect of sIL-13R α 2-Fc was unlikely to be explained by changes in Th1 or Th2-type cytokine expression. Therefore, in subsequent experiments, we investigated the patterns of collagen I (Col I) and collagen III (Col III) mRNA expression to determine whether the sIL-13R α 2-Fc-induced reduction in fibrosis was accompanied by direct changes in the expression of these two important collagen producing genes¹⁹. As shown in Figure 6, IL-13 blockade significantly reduced Col I and Col III mRNA expression in both WT and IL-4-deficient mice. There was no change in the infection-induced levels Col I or Col III mRNAs in IL-4-deficient mice and when compared with sIL-13R α 2-Fc-treated WT mice, there was no further reduction in similarly treated IL-4-deficient mice.

IL-13 stimulates collagen production in mouse 3T3 fibroblasts.

Having shown that IL-13 blockade *in vivo* significantly reduced Col I and Col III mRNA expression in the liver of infected WT and IL-4-deficient mice, we wanted to determine whether IL-13 would directly stimulate collagen synthesis in fibroblasts. To answer this question, we examined the induction of type I collagens in murine 3T3 fibroblasts by Western blotting. As shown in Fig. 7, IL-13 induced collagen synthesis 48 h after stimulation. Minimal type I collagen was detected in unstimulated cells (Fig 7, lane 1) or at earlier time points in the cytokine-activated cultures (data not shown). IL-4 also induced collagen I synthesis (lane 2) and high levels of secreted collagen were easily detectable in the supernatants obtained from both cytokine-stimulated cultures (data not shown). The specificity of the reaction was confirmed by using purified collagen type I (lane 5) and bacterial collagenase treatments showed that the antibodies were specific for collagen (data not shown).

DISCUSSION

A CD4⁺ Th2-type cytokine pattern dominates the immune response in mice infected with *S. mansoni*^{12,13}. Previous IL-4 depletion studies and experiments with IL-4-deficient mice however, failed to show an indispensable role for this cytokine in the pathogenesis of schistosomiasis^{15,23,24}. Indeed, while a partial reduction in fibrosis was observed in some studies¹⁵, egg-induced granuloma formation could proceed in the complete absence of IL-4^{23,24}. In contrast to these observations, granuloma formation

and the development of hepatic fibrosis was severely impaired in Stat6-deficient mice¹⁶, which display a major defect in the production of several Th2-associated cytokines⁴⁶. IL-4 and IL-13 both signal through Stat6, therefore the apparent differences in pathology observed between infected IL-4-deficient and Stat6-deficient mice may be explained by IL-13. Nevertheless, the distinct contributions of IL-4 and IL-13 in disease progression can not be discerned from studies in Stat6 or IL-4-deficient mice alone. In this study, we used a potent inhibitor of IL-13 in infected WT and IL-4-deficient mice and demonstrate that IL-13 and IL-4 exhibit redundant, as well as unique roles in the pathogenesis of schistosomiasis.

Several studies have shown that Th2-type cytokine responses can develop *in vivo* in the absence of IL-4 or the IL-4 receptor^{26,39}, which is consistent with our findings since reduced but significant IL-13, IL-10, and IL-5 expression was detected in the mesenteric lymph nodes (Fig. 4) and livers (Fig. 5) of infected IL-4-deficient mice. Their production was also highly dependent on a CD4⁺ T cell response (Fig. 4), further indicating that a conventional Th2-type response was established. These findings provide evidence that while maximal IL-13 expression is dependent on IL-4, the continued production of IL-13 might explain the maintenance of a significant granulomatous response in the absence of IL-4²³⁻²⁵. Indeed, while blocking IL-13 alone had no effect on granuloma size in WT mice, inhibiting the residual IL-13 in IL-4-deficient mice resulted in a marked and highly significant reduction in granuloma volume (Fig. 2A). These findings demonstrate that IL-4 and IL-13 are both sufficient to mediate granuloma development, and formally explain the production of granulomas in IL-4-deficient mice versus the nearly complete lack of granulomas in Stat6-deficient mice^{16,24}. They also support recent findings in the pulmonary egg granuloma model³⁰. Because granulomas serve an important host-protective role by walling off potentially lethal hepatotoxins released by the eggs⁴⁷, the host may have evolved redundant mechanisms for granuloma formation in order to ensure a favorable host-parasite relationship.

While these observations clearly demonstrate that IL-4 and IL-13 actively participate in granuloma formation, unique roles for both cytokines in mast cell recruitment, tissue eosinophilia, and most importantly, the generation of hepatic fibrosis were revealed in these studies. Histological examinations of liver sections from infected mice demonstrated that IL-13 is not required for mast cell (Table III) or eosinophil (Fig.

2B) differentiation and recruitment, since granulomas of sIL-13R α 2-Fc-treated WT mice showed no decrease in either cell type. In fact, eosinophil numbers were significantly increased in the lesions of IL-13-inhibited WT mice (Fig. 2B), suggesting that IL-13 may partially antagonize this effect. In contrast, mast cells were almost completely absent from the lesions in IL-4-deficient mice and eosinophils were decreased by over 50%. Interestingly, IL-13 appears to partially support the reduced but significant egg-induced tissue eosinophilia in IL-4-deficient mice since eosinophils were reduced to below 10% in the IL-4-deficient/sIL-13R α 2-Fc-treated animals. Nevertheless, these data indicate that IL-4 is the dominant cytokine responsible for the development of eosinophil and mast cell populations within granulomas.

Probably the most important advance from this study was the finding that hepatic fibrosis could be blocked by sIL-13R α 2-Fc. Indeed, microscopic (Table III), biochemical (Fig. 2C), and molecular techniques (Fig. 6) all indicated that IL-13, not IL-4, plays the major role in the development of egg-induced liver fibrosis. Previous studies showed that the Th1/Th2 cytokine balance can significantly effect the extent of tissue fibrosis in *S. mansoni* infected mice¹⁹. Nevertheless, this study suggests that the effects of sIL-13R α 2-Fc were not mediated through a skewing of the Th cell cytokine response. Blocking IL-13 had no significant effect on the production of IFN- γ , IL-4, IL-5, IL-10, or IL-13 by mesenteric lymph node (Fig. 4) or spleen cells *in vitro* and there was also no change in cytokine mRNA expression *in vivo*, at the site of lesion formation (Fig. 5). In contrast to these observations, IL-4-deficient mice displayed an increased IFN- γ response in the draining lymph nodes (Fig. 4) and decreased IL-5 and IL-13 expression in both the lymph nodes (Fig. 4) and liver (Fig. 5). Thus, the slight reduction in fibrosis detected in IL-4-deficient mice by hydroxyproline analysis (Fig. 2C) may be attributable to decreased IL-13 production. The fact that IL-4 production was unaffected by IL-13 blockade, yet fibrosis was maximally reduced in these animals emphasizes the important role played by IL-13. Indeed, sIL-13R α 2-Fc-treated IL-4-deficient mice showed little additional decrease in hydroxyproline levels (Fig. 2C) and no difference in Collagen I or III mRNA expression (Fig. 6) over that observed in similarly-treated WT mice. There was also no change in Collagen I or III mRNA expression in control-Fc-treated IL-4-deficient mice when compared with WT animals, further de-emphasizing the contribution of IL-4. Moreover, *in vitro* studies with 3T3 cells demonstrated for the first time the ability of IL-13 to

stimulate collagen production in fibroblasts (Fig. 7), thus the effects of IL-13 on fibrosis may be more direct and not dependent upon modulations in the Th1/Th2 cytokine response. In support of this conclusion, recent studies demonstrated that IL-13 receptors are expressed on fibroblasts³² and that IL-13 increases adhesion molecule and inflammatory cytokine expression in human lung fibroblasts⁴⁸. Finally, although IL-13 (Fig. 7) and IL-4⁴⁹ are both capable of promoting collagen production in fibroblasts, the fact that cultured lymph node cells produced nearly 100-fold more IL-13 than IL-4 (Fig. 4), only serves to emphasize the potentially important contribution of IL-13 in this process. Indeed, studies in the pulmonary granuloma model revealed that IL-4 mRNA expression is more tightly regulated at the site of lesion formation, while the induction of IL-13 mRNA is much more sustained over time³⁰. Nevertheless, we have not examined the kinetics of IL-4 and IL-13 mRNA expression in infected animals, so we can not say whether a similar pattern holds in the granulomatous livers.

IL-13 was also recently shown to be important for resistance against intestinal nematodes^{27,37-39}. Studies in IL-4³⁹ and IL-13-deficient mice^{37,38} suggested that IL-13, in contrast to IL-4, plays a requisite role in expulsion of both *N. brasiliensis* and *T. muris*. Nevertheless, the specific mechanism of worm expulsion remains unknown, although IL-4 and IL-13-induced changes in epithelial cells and gut physiology have been suggested as possible targets^{50,51}. IL-13 also plays a central role in murine asthma models. In these studies, IL-13 was found to be necessary and sufficient for the expression of allergic asthma^{40,41}. Subepithelial fibrosis and airway smooth muscle hypertrophy are common features of chronic severe asthma⁵ and chronic pulmonary fibrosis is associated with the production of type III and type I collagen in the early and late stages of the disease, respectively. Thus the link between IL-13 and fibrosis revealed in our study elucidates the etiology of several important human diseases and provides more effective modes of treatment of fibrotic diseases in general.

Our previous studies showed that an egg specific IL-12-induced Th1 memory response could effectively reduce hepatic fibrosis in subsequently-infected mice¹⁹. The reduction in pathology was accompanied by a switch in the normal Th2 response to one dominated by Th1-type cytokines. Findings from the current study suggest that the anti-pathology effects of this IL-12 -based vaccination protocol may be explained by the inhibition of IL-13. Interestingly, a second study using a different protocol showed that

repeated rIL-12 injections given at 6 to 8 weeks, during the Th2-dominated phase of granuloma development, was almost completely ineffective at blocking granuloma formation and fibrosis⁵². Related studies have suggested that IL-12 is less capable of modulating established Th2-type responses⁵³, which likely explains the failure to
5 modulate pathology in the latter study⁵². In contrast to these findings, sIL-13R α 2-Fc was extremely effective at reducing hepatic fibrosis, even though administered only during the later stages of infection. These findings indicate that IL-13 antagonism is a much more effective therapeutic approach to reduce fibrosis in situations where pathogenic Th2-type immune responses have already been established. In summary, our findings provide
10 evidence that IL-13 inhibitors, such as the sIL-13R α 2-Fc, are of general therapeutic benefit in preventing fibrosis associated with chronic infectious disease and demonstrate the important and non-redundant role of IL-13 in the pathogenesis of schistosomiasis.

METHODS

15 *Animals, Parasites and Ag preparations*

6-8 week old female C57BL/6 and IL-4-deficient mice (C57BL/6 background, 10th backcross) were obtained from Taconic Farms, Inc. (Germantown, NY). All mice were housed in a NIH American Association for the Accreditation of Laboratory Animal Care-approved animal facility in sterile filter-top cages and maintained on sterile water. Cercariae of a Puerto
20 Rican strain of *Schistosoma mansoni* (NMRI) were obtained from infected *Biomphalaria glabrata* snails (Biomedical Research Institute, Rockville, MD). Soluble egg antigen (SEA) was purified from homogenized eggs, as previously described¹⁵.

Reagents

25 The soluble IL-13 receptor α 2-Fc fusion protein (sIL-13R α 2-Fc) was prepared as previously described³⁵ and provided by Genetics Institute, Cambridge MA. Endotoxin contamination was <2 EU/mg, as determined with the Cape Cod Associates LAL assay (Limulus Amebocyte Lysate, Woods Hole, MA). The in vitro ID50, as determined by the ability to neutralize 3ng/ml of murine IL-13 in the B9 proliferation assay, was
30 approximately 10 ng/ml. Human IgG (control-Fc), which was used as a control for sIL-13R α 2-Fc, was affinity purified by recombinant Protein A-Sepharose chromatography, as

described for sIL-13R α 2-Fc³⁵. As described previously, the control-Fc had no detectable effect on pathology or cytokine expression in infected mice³⁰.

Infection and treatments

5 Mice were infected by percutaneous challenge of tail skin for 40 min in water containing between 20 and 25 cercariae. Animals were treated with either a human control-Fc or with the sIL-13R α 2-Fc by i.p. injection in 0.5 ml PBS, every other day after the onset of egg production (week 5). The optimal concentration for in vivo use (200 μ g/mouse/day) was chosen based on kinetic assays and on dose response experiments in
10 sensitized/i.v. egg-injected mice³⁰. Sera were collected from mice on the day of sacrifice. All animals were sacrificed by i.p. administration of sodium pentobarbital (18 mg/mouse, Sigma, St. Louis, MO) on week 8 and perfused with citrated saline to assess worm burdens¹⁵. No mortality was observed among any of the treated groups.

Histopathology and fibrosis measurement

15 For measurement of granulomas, approximately half of the liver was fixed with Bouin-Hollande fixative and processed as previously described¹⁵. The size of hepatic granulomas was determined in histological sections stained by Wright's Giemsa stain (Histopath of America, Clinton, MD). The diameters of each granuloma containing a
20 single viable egg were measured with an ocular micrometer and the volume of each granuloma calculated assuming a spherical shape. The mean of the longest diameter and the diameter perpendicular to that was used. The percentage of eosinophils, mast cells and other cell types were evaluated in the same sections. Parenchymal necrosis was scored on a scale of 0-4, with 0 being the least and 4 being the most extensive necrosis. The
25 frequency of mast cells was also assigned on a similar scale, using a range from 0-4. The number of schistosome eggs in the liver and gut and the collagen content of the liver, determined as hydroxyproline, were measured as described previously¹⁵. Fibrosis was also scored histologically using sections stained with picrosirius red. The picrosirius reagent stains collagen specifically and when sections are viewed under polarizing light,
30 the bright areas where collagen is deposited are illuminated. All granulomas within each section were scored for picrosirius (red) "density" based on a scale 1-4, and a second measurement of "area involved" was also determined using the same scale. The total

fibrosis score was determined by multiplying the density and area for each granuloma (ie. a score of 16 would be the maximum). An average of 30 granulomas per mouse was included in all analyses. To control for consistency, the same individual scored all histological features and had no knowledge of the experimental design.

5

Isolation and purification of RNA

Two portions of the liver from each animal were combined and placed in 1 ml of RNA-STAT 60 (Tel-Test), frozen on dry ice and kept at -70°C until use. Tissues were homogenized using a tissue polytron (Omni International Inc., Waterbury, CT) and total
10 RNA was extracted following the recommendations of the manufacturer. The RNA was resuspended in DEPC-treated water and quantitated spectrophotometrically.

RT-PCR detection of cytokine mRNA

A RT-PCR procedure was used to determine relative quantities of mRNA for IL-4,
15 IL-5, IL-10, IL-13, IFN- γ , collagen I, collagen III, TGF β 1, TGF β 2, and HPRT (hypoxanthine-guanine phosphoribosyl transferase). The cDNA was obtained after reverse transcription of 1 μ g of RNA as described ¹⁴. The primers and probes for all genes were previously published ^{14,19,54}. The PCR cycles used for each cytokine were as follows: IL-4 (33), IL-5 (31), IFN- γ (29), collagen I (26), collagen III (22), TGF β 1 (34), TGF β 2
20 (34), and HPRT (23).

Analysis and quantification of PCR products

The amplified DNA was analyzed by electrophoresis, Southern blotting and hybridization with non-radioactive cytokine-specific probes as previously described ¹⁴.
25 The PCR products were detected using a ECL detection system (Amersham). The chemiluminescent signals were quantified using a flat-bed scanner (Microtek model 600 ZS, Torrance, CA). The amount of PCR product was determined by comparing the ratio of cytokine-specific signal density to that of HPRT-specific signal density for individual samples. Arbitrary densitometric units for individual samples were subsequently multiplied
30 by a factor of 100.

In vitro cultures

Mesenteric lymph node (MLN) cells and spleens were extracted from the mice and single cell suspensions were prepared. Red blood cells were lysed by osmotic treatment with ACK lysing buffer (Biofluids, Inc., Rockville, MD). Cells were placed in RPMI 1640 medium supplemented with 10% FCS, 2mM glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin, 25 mM HEPES, 1mM sodium pyruvate, 0.1 mM nonessential amino acids, and 50 µM 2-ME at 37°C in 5% CO₂. Cells were plated in 24-well plates (3 x 10⁶/ml, 1ml) and stimulated with SEA (20 µg/ml) and supernatants were collected after 72 h to measure the levels of IL-4, IL-5, IL-10, IL-13 and IFN-γ. Additional SEA-stimulated cultures were also treated with 50 µg/ml of anti-CD4 mAb (GK1.5). Cultures treated with anti-CD4 mAb alone showed no change in cytokine expression when compared with that observed in medium control cultures (data not shown). IL-5, IL-10, and IFN-γ were measured using specific sandwich ELISA¹⁵. IL-13 levels were measured using murine IL-13 ELISA kits (R&D Systems, Minneapolis, MN). Cytokine levels were calculated from curves prepared with recombinant cytokines. IL-4 was measured using the IL-4 sensitive cell line CT.4S. Proliferation of these cells was quantified by (³H)TdR incorporation, and the amount of cytokine was determined by comparison with known amounts of recombinant IL-4.

Western blot detection of collagen I

3T3 fibroblasts were cultured in RPMI 1640 medium supplemented with 10% FCS, 2mM glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin, 25 mM HEPES, 1mM sodium pyruvate, 0.1 mM nonessential amino acids, and 50 µM 2-ME at 37°C in 5% CO₂. Confluent cells were plated in 24-well plates (500,000 cells/ml) and stimulated with IL-4 (1000 U/ml) or rIL-13 (R&D Systems, Minneapolis, MN) (20 ng/ml) for 6, 24 and 48 hs. Culture supernatants were collected to analyze secreted collagen I. Cells were washed once with phosphate buffered saline and lysed with SDS-PAGE sample buffer. Cell lysates and culture supernatants were submitted to electrophoretic separation in 6% tris-glycine gels (Novel Experimental Technology, San Diego, CA) using reducing conditions, and transferred to nitrocellulose membranes (Schleicher & Schuell, Keene, NH). Blots were probed with rabbit IgG anti-mouse type I collagen (Biodesign International, Kennebunk, ME) and peroxidase labelled anti-rabbit IgG (Amersham Pharmacia Biotech,

Inc., Piscataway, NJ) was used as a second Ab. The bands were visualized using a western blot chemiluminescence reagent (NEN Life Science Products, Boston, MA). To confirm identity of the collagen bands, cell lysates were treated with 0.5 mg/ml of collagenase (Boehringer Mannheim, Indianapolis, IN) in PBS, supplemented with 1 mM CaCl₂ and 1% FCS, for 1 h at 37°C. A purified rat collagen I preparation was also used as a control.

Statistics

Schistosome worm and egg numbers, changes in cytokine mRNA, and values for secreted cytokine proteins were compared using Student's two-tailed t test. Hepatic fibrosis was compared by analysis of covariance, using the log of total liver eggs as the covariate and the log of hydroxyproline per egg. $p < 0.05$ was considered significant.

References

1. Rosenstein, B.J. & Zeitlin, P.L. Cystic fibrosis. *Lancet* **351**, 277-282 (1998).
2. Lee, J.K. *et al.* A serine elastase inhibitor reduces inflammation and fibrosis and preserves cardiac function after experimentally-induced murine myocarditis. *Nat Med* **4**, 1383-1391 (1998).
3. Lahita, R.G. Collagen disease: the enemy within. *Int J Fertil Womens Med* **43**, 229-234 (1998).
4. Kuroda, K., Tsukifuji, R. & Shinkai, H. Increased expression of heat-shock protein 47 is associated with overproduction of type I procollagen in systemic sclerosis skin fibroblasts. *J Invest Dermatol* **111**, 1023-1028 (1998).
5. Bento, A.M. & Hershenson, M.B. Airway remodeling: potential contributions of subepithelial fibrosis and airway smooth muscle hypertrophy/hyperplasia to airway narrowing in asthma. *Allergy Asthma Proc* **19**, 353-358 (1998).
6. Wahl, S.M. *et al.* Bacterial cell wall-induced hepatic granulomas. An in vivo model of T cell-dependent fibrosis. *J Exp Med* **163**, 884-902 (1986).
7. Henderson, G.S. *et al.* Two distinct pathological syndromes in male CBA/J inbred mice with chronic *Schistosoma mansoni* infections. *Am J Pathol* **142**, 703-714 (1993).

8. Trojanowska, M., LeRoy, E.C., Eckes, B. & Krieg, T. Pathogenesis of fibrosis: type 1 collagen and the skin. *J Mol Med* **76**, 266-274 (1998).
9. Johnson, L.L., Dyer, R. & Hupe, D.J. Matrix metalloproteinases. *Curr Opin Chem Biol* **2**, 466-471 (1998).
- 5 10. Cheever, A.W. & Yap, G.S. Immunologic basis of disease and disease regulation in schistosomiasis. *Chem Immunol* **66**, 159-176 (1997).
11. Bergquist, N.R. Schistosomiasis vaccine development: progress and prospects [In Process Citation]. *Mem Inst Oswaldo Cruz* **93**, 95-101 (1998).
12. Grzych, J.M. *et al.* Egg deposition is the major stimulus for the production of Th2
10 cytokines in murine schistosomiasis mansoni. *J.Immunol.* **146**, 1322-1327 (1991).
13. Pearce, E.J., Caspar, P., Grzych, J.M., Lewis, F.A. & Sher, A. Downregulation of Th1 cytokine production accompanies induction of Th2 responses by a parasitic helminth, *Schistosoma mansoni*. *J.Exp.Med.* **173**, 159-162 (1992).
14. Wynn, T.A. *et al.* Analysis of cytokine mRNA expression during primary
15 granuloma formation induced by eggs of *Schistosoma mansoni*. *J.Immunol.* **151**, 1430-1440 (1993).
15. Cheever, A.W. *et al.* Anti-IL-4 treatment of *Schistosoma mansoni*-infected mice inhibits development of T cells and non-B, non-T cells expressing Th2 cytokines while decreasing egg-induced hepatic fibrosis. *J.Immunol.* **153**, 753-754 (1994).
- 20 16. Kaplan, M.H., Whitfield, J.R., Boros, D.L. & Grusby, M.J. Th2 cells are required for the *Schistosoma mansoni* egg-induced granulomatous response. *J Immunol* **160**, 1850-1856 (1998).
17. Chensue, S.W. *et al.* Role of interleukin-4 and gamma-interferon in *Schistosoma mansoni* egg-induced hypersensitivity granuloma formation: orchestration,
25 relative contribution and relationship to macrophage function. *J.Immunol.* **148**, 900-910 (1992).
18. Henderson, G.S., Lu, X., McCurley, T.L. & Colley, D.G. In vivo molecular analysis of lymphokines involved in the murine immune response during *Schistosoma mansoni* infection: II. Quantitation of IL-4 mRNA, IFN-g mRNA,
30 and IL-2 mRNA levels in the granulomatous livers, mesenteric lymph nodes, and spleens during the course of modulation. *J.Immunol.* **148**, 2261-2267 (1992).

19. Wynn, T.A. *et al.* An IL-12-based vaccination method for preventing fibrosis induced by schistosome infection. *Nature* **376**, 594-596 (1995).
20. Secor, W.E., Stewart, S.J. & Colley, D.G. Eosinophils and immune mechanisms. VI. The synergistic combination of granulocyte-macrophage colony-stimulating factor and IL-5 accounts for eosinophil-stimulation promoter activity in *Schistosoma mansoni*-infected mice. *J Immunol* **144**, 1484-1489 (1990).
21. Le Gros, G., Ben-Sasson, S.Z., Seder, R., Finkelman, F.D. & Paul, W.E. Generation of interleukin 4 (IL-4)-producing cells in vivo and in vitro: IL-2 and IL-4 are required for in vitro generation of IL-4-producing cells. *J Exp Med* **172**, 921-929 (1990).
22. Kopf, M. *et al.* Disruption of the murine IL-4 gene blocks Th2 cytokine responses. *Nature* **362**, 245-248 (1993).
23. Metwali, A. *et al.* The granulomatous response in murine Schistosomiasis mansoni does not switch to Th1 in IL-4-deficient C57BL/6 mice. *J Immunol* **157**, 4546-4553 (1996).
24. Pearce, E.J. *et al.* *Schistosoma mansoni* in IL-4-deficient mice. *Int Immunol* **8**, 435-444 (1996).
25. Chensue, S.W., Warmington, K., Ruth, J.H., Lukacs, N. & Kunkel, S.L. Mycobacterial and schistosomal antigen-elicited granuloma formation in IFN-gamma and IL-4 knockout mice: analysis of local and regional cytokine and chemokine networks. *J Immunol* **159**, 3565-3573 (1997).
26. Noben-Trauth, N. *et al.* An interleukin 4 (IL-4)-independent pathway for CD4+ T cell IL-4 production is revealed in IL-4 receptor-deficient mice. *Proc Natl Acad Sci U S A* **94**, 10838-10843 (1997).
27. Barner, M., Mohrs, M., Brombacher, F. & Kopf, M. Differences between IL-4R alpha-deficient and IL-4-deficient mice reveal a role for IL-13 in the regulation of Th2 responses. *Curr Biol* **8**, 669-672 (1998).
28. Amiri, P. *et al.* Tumour necrosis factor alpha restores granulomas and induces parasite egg-laying in schistosome-infected SCID mice. *Nature* **356**, 604-607 (1992).

29. Hernandez, H.J., Wang, Y. & Stadecker, M.J. In infection with *Schistosoma mansoni*, B cells are required for T helper type 2 cell responses but not for granuloma formation. *J Immunol* **158**, 4832-4837 (1997).
30. Chiaramonte, M.G. *et al.* IL-13 is a key regulatory cytokine for Th2 cell-mediated pulmonary granuloma formation and IgE responses induced by *Schistosoma mansoni* eggs. *J Immunol* **162**, 920-930 (1999).
31. de Vries, J.E. The role of IL-13 and its receptor in allergy and inflammatory responses. *J Allergy Clin Immunol* **102**, 165-169 (1998).
32. Murata, T., Husain, S.R., Mohri, H. & Puri, R.K. Two different IL-13 receptor chains are expressed in normal human skin fibroblasts, and IL-4 and IL-13 mediate signal transduction through a common pathway. *Int Immunol* **10**, 1103-1110 (1998).
33. Emson, C.L., Bell, S.E., Jones, A., Wisden, W. & McKenzie, A.N. Interleukin (IL)-4-independent induction of immunoglobulin (Ig)E, and perturbation of T cell development in transgenic mice expressing IL-13. *J Exp Med* **188**, 399-404 (1998).
34. McKenzie, G.J. *et al.* Impaired development of Th2 cells in IL-13-deficient mice. *Immunity* **9**, 423-432 (1998).
35. Donaldson, D.D. *et al.* The Murine IL-13Ra2: Molecular Cloning, Characterization and Comparison with Murine IL-13Ra1. *J. Immunol.* **161**, 2317-24. (1998).
36. Minty, A. *et al.* Interleukin-13 is a new human lymphokine regulating inflammatory and immune responses. *Nature* **362**, 248-250 (1993).
37. McKenzie, G.J., Bancroft, A., Grencis, R.K. & McKenzie, A.N. A distinct role for interleukin-13 in Th2-cell-mediated immune responses. *Curr Biol* **8**, 339-342 (1998).
38. Bancroft, A.J., McKenzie, A.N. & Grencis, R.K. A critical role for IL-13 in resistance to intestinal nematode infection. *J Immunol* **160**, 3453-3461 (1998).
39. Urban, J.F., Jr. *et al.* IL-13, IL-4R α , and Stat6 are required for the expulsion of the gastrointestinal nematode parasite *Nippostrongylus brasiliensis*. *Immunity* **8**, 255-264 (1998).

40. Wills-Karp, M. *et al.* Interleukin-13: central mediator of allergic asthma. *Science* **282**, 2258-2261 (1998).
41. Grunig, G. *et al.* Requirement for IL-13 independently of IL-4 in experimental asthma. *Science* **282**, 2261-2263 (1998).
- 5 42. Hilton, D.J. *et al.* Cloning and characterization of a binding subunit of the interleukin 13 receptor that is also a component of the interleukin 4 receptor. *Proc Natl Acad Sci U S A* **93**, 497-501 (1996).
43. Miloux, B. *et al.* Cloning of the human IL-13R alpha1 chain and reconstitution with the IL4R alpha of a functional IL-4/IL-13 receptor complex. *FEBS Lett* **401**, 163-166 (1997).
- 10 44. Gauchat, J.F. *et al.* A novel 4-kb interleukin-13 receptor alpha mRNA expressed in human B, T, and endothelial cells encoding an alternate type-II interleukin-4/interleukin-13 receptor. *Eur J Immunol* **27**, 971-978 (1997).
45. Warren, K.S. Schistosomiasis: host-pathogen biology. *Rev Infect Dis* **4**, 771-775 (1982).
- 15 46. Kaplan, M.H., Schindler, U., Smiley, S.T. & Grusby, M.J. Stat6 is required for mediating responses to IL-4 and for development of Th2 cells. *Immunity* **4**, 313-319 (1996).
47. Dunne, D.W., Jones, F.M. & Doenhoff, M.J. The purification, characterization, serological activity and hepatotoxic properties of two cationic glycoproteins (alpha 1 and omega 1) from *Schistosoma mansoni* eggs. *Parasitology* **103 Pt 2**, 225-236 (1991).
- 20 48. Doucet, C. *et al.* IL-4 and IL-13 specifically increase adhesion molecule and inflammatory cytokine expression in human lung fibroblasts. *Int Immunol* **10**, 1421-1433 (1998).
- 25 49. Serpier, H. *et al.* Antagonistic effects of interferon-gamma and interleukin-4 on fibroblast cultures. *J Invest Dermatol* **109**, 158-162 (1997).
50. Zund, G., Madara, J.L., Dzus, A.L., Awtrey, C.S. & Colgan, S.P. Interleukin-4 and interleukin-13 differentially regulate epithelial chloride secretion. *J Biol Chem* **271**, 7460-7464 (1996).
- 30

51. Finkelman, F.D. *et al.* Cytokine regulation of host defense against parasitic gastrointestinal nematodes: lessons from studies with rodent models. *Annu Rev Immunol* **15**, 505-533 (1997).
52. Boros, D.L. & Whitfield, J.R. Enhanced Th1 and dampened Th2 responses synergize To inhibit acute granulomatous and fibrotic responses in murine schistosomiasis mansoni [In Process Citation]. *Infect Immun* **67**, 1187-1193 (1999).
53. Wang, Z.E. *et al.* Interferon gamma-independent effects of interleukin 12 administered during acute or established infection due to *Leishmania major*. *Proc Natl Acad Sci U S A* **91**, 12932-12936 (1994).
54. Wynn, T.A., Eltoun, I., Oswald, I.P., Cheever, A.W. & Sher, A. Endogenous interleukin 12 (IL-12) regulates granuloma formation induced by eggs of *Schistosoma mansoni* and exogenous IL-12 both inhibits and prophylactically immunizes against egg pathology. *J.Exp.Med.* **179**, 1551-1561 (1994).

All patent and literature references cited herein are incorporated by reference as if fully set forth.

What is claimed is:

1. A method of treating tissue fibrosis in a mammalian subject, said method comprising administering a therapeutically effective amount of a pharmaceutical composition comprising a protein and a pharmaceutically acceptable carrier, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain..

2. The method of claim 1 wherein said tissue fibrosis affects a tissue selected from the group consisting of liver, skin epidermis, skin endodermis, muscle, tendon, cartilage, cardiac tissue, pancreatic tissue, lung tissue, uterine tissue, neural tissue, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract and gut.

3. The method of claim 2 wherein said tissue is liver.

4. The method of claim 2 wherein said fibrosis is that resulting from infection with schistosoma.

5. The method of claim 1 wherein said fibrosis is that resulting from healing of a wound.

6. A method of inhibiting formation of tissue fibrosis in a mammalian subject, said method comprising administering a therapeutically effective amount of a pharmaceutical composition comprising a protein and a pharmaceutically acceptable carrier, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain..

7. The method of claim 6 wherein said tissue fibrosis affects a tissue selected from the group consisting of liver, skin epidermis, skin endodermis, muscle, tendon, cartilage, cardiac tissue, pancreatic tissue, lung tissue, uterine tissue, neural tissue, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract and gut.

8. The method of claim 7 wherein said tissue is liver.

9. The method of claim 7 wherein said fibrosis is that resulting from infection with schistosoma.

10. The method of claim 6 wherein said fibrosis is that resulting from healing of a wound.

11. The method of claim 10 wherein said wound is a surgical incision.

12. The method of claim 5 wherein said wound is a surgical incision.

13. A method of treating tissue fibrosis in a mammalian subject, said method comprising administering a therapeutically effective amount of a composition comprising (a) a molecule selected from the group consisting of an IL-13 antagonist and an IL-4 antagonist, and (b) a pharmaceutically acceptable carrier.

14. The method of claim 13 wherein said antagonist is selected from the group consisting of an IL-13bc protein, a soluble form of IL-13R α 1, an antibody to IL-13 or an IL-13-binding fragment thereof, an antibody to IL-13bc or an IL-13bc-binding fragment thereof, an antibody to IL-13R α 1 or an IL-13R α 1-binding fragment thereof, IL-13R-binding mutants of IL-4, a small molecule capable of inhibiting the interaction of IL-13 with IL-13bc and a small molecule capable of inhibiting the interaction of IL-13 with IL-13R α 1.

15. The method of claim 14 wherein said IL-13bc protein is a protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
- (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
- (d) the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
- (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and
- (g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain.

16. The method of claim 13 wherein said tissue fibrosis affects a tissue selected from the group consisting of liver, skin epidermis, skin endodermis, muscle, tendon, cartilage, cardiac tissue, pancreatic tissue, lung tissue, uterine tissue, neural tissue, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract and gut.

17. The method of claim 16 wherein said tissue is liver.

18. The method of claim 17 wherein said fibrosis is that resulting from infection with schistosoma.

19. The method of claim 13 wherein said fibrosis is that resulting from healing of a wound.

20. The method of claim 19 wherein said wound is a surgical incision.
21. A method of inhibiting formation of tissue fibrosis in a mammalian subject, said method comprising administering a therapeutically effective amount of a composition comprising (a) a molecule selected from the group consisting of an IL-13 antagonist and an IL-4 antagonist, and (b) a pharmaceutically acceptable carrier.
22. The method of claim 21 wherein said antagonist is selected from the group consisting of an IL-13bc protein, a soluble form of IL-13R α 1, an antibody to IL-13 or an IL-13-binding fragment thereof, an antibody to IL-13bc or an IL-13bc-binding fragment thereof, an antibody to IL-13R α 1 or an IL-13R α 1-binding fragment thereof, IL-13R-binding mutants of IL-4, a small molecule capable of inhibiting the interaction of IL-13 with IL-13bc and a small molecule capable of inhibiting the interaction of IL-13 with IL-13R α 1.
23. The method of claim 22 wherein said IL-13bc protein is a protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:2;
 - (b) the amino acid sequence of SEQ ID NO:2 from amino acids 22 to 334;
 - (c) the amino acid sequence of SEQ ID NO:2 from amino acids 357 to 383;
 - (d) the amino acid sequence of SEQ ID NO:4;
 - (e) the amino acid sequence of SEQ ID NO:4 from amino acids 26 to 341;
 - (f) the amino acid sequence of SEQ ID NO:4 from amino acids 363 to 380; and

(g) fragments of (a)-(f) having a biological activity of the IL-13 receptor binding chain.

24. The method of claim 21 wherein said tissue fibrosis affects a tissue selected from the group consisting of liver, skin epidermis, skin endodermis, muscle, tendon, cartilage, cardiac tissue, pancreatic tissue, lung tissue, uterine tissue, neural tissue, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract and gut.

25. The method of claim 24 wherein said tissue is liver.

26. The method of claim 25 wherein said fibrosis is that resulting from infection with schistosoma.

27. The method of claim 21 wherein said fibrosis is that resulting from healing of a wound.

28. The method of claim 27 wherein said wound is a surgical incision.

29. The method of claim 21 wherein said antagonist is selected from the group consisting of a soluble form of IL-4R, an antibody to IL-4 or an IL-4-binding fragment thereof, an antibody to IL-4R or an IL-4R-binding fragment thereof, and a small molecule capable of inhibiting the interaction of IL-4 with IL-4R.

30. The method of claim 13 wherein said antagonist is selected from the group consisting of a soluble form of IL-4R, an antibody to IL-4 or an IL-4-binding fragment thereof, an antibody to IL-4R or an IL-4R-binding fragment thereof, and a small molecule capable of inhibiting the interaction of IL-4 with IL-4R.

Figure 1




Figure 2




Figure 3




Figure 4A




Figure 4B




Figure 4C




Figure 5




Figure 6





Figure 7



SEQUENCE LISTING

<110> Wynn, Thomas A.
Chiaramonte, Monica G.
Collins, Mary
Donaldson, Debra
Fitz, Lori
Neben, Tamlyn
Whitters, Matthew
Wood, Clive
Genetics Institute, Inc.

<120> TREATMENT OF FIBROSIS BY ANTAGONISM OF IL-13 AND IL-13
RECEPTOR CHAINS

<130> GI 5268A2A

<140>

<141>

<160> 9

<170> PatentIn Ver. 2.0

<210> 1

<211> 1525

<212> DNA

<213> Mus sp.

<220>

<221> CDS

<222> (256)..(1404)

<400> 1

gaattcggca cgagggagag gaggagggaa agatagaaag agagagagaa agattgcttg 60

ctaccctga acagtgcact ctctcaagac agtgctttgc tcttcacgta taaggaagga 120

aaacagtaga gattcaattt agtgtctaata gtggaaagga ggacaaagag gtcttgtgat 180

aactgcctgt gataatacat ttcttgagaa accatattat tgagtagagc tttcagcaca 240

ctaaatcctg gagaa atg gct ttt gtg cat atc aga tgc ttg tgt ttc att 291

Met Ala Phe Val His Ile Arg Cys Leu Cys Phe Ile

1 5 10

ctt ctt tgt aca ata act ggc tat tct ttg gag ata aaa gtt aat cct 339

Leu Leu Cys Thr Ile Thr Gly Tyr Ser Leu Glu Ile Lys Val Asn Pro

15 20 25

cct cag gat ttt gaa ata ttg gat cct gga tta ctt ggt tat ctc tat 387

Pro Gln Asp Phe Glu Ile Leu Asp Pro Gly Leu Leu Gly Tyr Leu Tyr

30 35 40

ttg caa tgg aaa cct cct gtg gtt ata gaa aaa ttt aag ggc tgt aca 435

Leu Gln Trp Lys Pro Pro Val Val Ile Glu Lys Phe Lys Gly Cys Thr

45 50 55 60

cta gaa tat gag tta aaa tac cga aat gtt gat agc gac agc tgg aag 483

Leu Glu Tyr Glu Leu Lys Tyr Arg Asn Val Asp Ser Asp Ser Trp Lys

"Express Mail" mailing label number EL56479602005

Date of Deposit 21 June 2000

I hereby certify that this paper or fee is being
deposited with the United States Postal Service
"Express Mail Post Office to Addressee" service
under 37 CFR 1.10 on the date indicated above
and is addressed to the Assistant Commissioner
For Patents, Washington, D.C. 20231

Mary H. Hanna

65	70	75	
act ata att act agg aat cta att tac aag gat ggg ttt gat ctt aat			531
Thr Ile Ile Thr Arg Asn Leu Ile Tyr Lys Asp Gly Phe Asp Leu Asn			
80	85	90	
aaa ggc att gaa gga aag ata cgt acg cat ttg tca gag cat tgt aca			579
Lys Gly Ile Glu Gly Lys Ile Arg Thr His Leu Ser Glu His Cys Thr			
95	100	105	
aat gga tca gaa gta caa agt cca tgg ata gaa gct tct tat ggg ata			627
Asn Gly Ser Glu Val Gln Ser Pro Trp Ile Glu Ala Ser Tyr Gly Ile			
110	115	120	
tca gat gaa gga agt ttg gaa act aaa att cag gac atg aag tgt ata			675
Ser Asp Glu Gly Ser Leu Glu Thr Lys Ile Gln Asp Met Lys Cys Ile			
125	130	135	140
tat tat aac tgg cag tat ttg gtc tgc tct tgg aaa cct ggc aag aca			723
Tyr Tyr Asn Trp Gln Tyr Leu Val Cys Ser Trp Lys Pro Gly Lys Thr			
145	150	155	
gta tat tct gat acc aac tat acc atg ttt ttc tgg tat gag ggc ttg			771
Val Tyr Ser Asp Thr Asn Tyr Thr Met Phe Phe Trp Tyr Glu Gly Leu			
160	165	170	
gat cat gcc tta cag tgt gct gat tac ctc cag cat gat gaa aaa aat			819
Asp His Ala Leu Gln Cys Ala Asp Tyr Leu Gln His Asp Glu Lys Asn			
175	180	185	
gtt gga tgc aaa ctg tcc aac ttg gac tca tca gac tat aaa gat ttt			867
Val Gly Cys Lys Leu Ser Asn Leu Asp Ser Ser Asp Tyr Lys Asp Phe			
190	195	200	
ttt atc tgt gtt aat gga tct tca aag ttg gaa ccc atc aga tcc agc			915
Phe Ile Cys Val Asn Gly Ser Ser Lys Leu Glu Pro Ile Arg Ser Ser			
205	210	215	220
tat aca gtt ttt caa ctt caa aat ata gtt aaa cca ttg cca cca gaa			963
Tyr Thr Val Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro Pro Glu			
225	230	235	
ttc ctt cat att agt gtg gag aat tcc att gat att aga atg aaa tgg			1011
Phe Leu His Ile Ser Val Glu Asn Ser Ile Asp Ile Arg Met Lys Trp			
240	245	250	
agc aca cct gga gga ccc att cca cca agg tgt tac act tat gaa att			1059
Ser Thr Pro Gly Gly Pro Ile Pro Pro Arg Cys Tyr Thr Tyr Glu Ile			
255	260	265	
gtg atc cga gaa gac gat att tcc tgg gag tct gcc aca gac aaa aac			1107
Val Ile Arg Glu Asp Asp Ile Ser Trp Glu Ser Ala Thr Asp Lys Asn			
270	275	280	
gat atg aag ttg aag agg aga gca aat gaa agt gaa gac cta tgc ttt			1155
Asp Met Lys Leu Lys Arg Arg Ala Asn Glu Ser Glu Asp Leu Cys Phe			
285	290	295	300
ttt gta aga tgt aag gtc aat ata tat tgt gca gat gat gga att tgg			1203
Phe Val Arg Cys Lys Val Asn Ile Tyr Cys Ala Asp Asp Gly Ile Trp			

```

                305                310                315
agc gaa tgg agt gaa gag gaa tgt tgg gaa ggt tac aca ggg cca gac 1251
Ser Glu Trp Ser Glu Glu Glu Cys Trp Glu Gly Tyr Thr Gly Pro Asp
                320                325                330

tca aag att att ttc ata gta cca gtt tgt ctt ttc ttt ata ttc ctt 1299
Ser Lys Ile Ile Phe Ile Val Pro Val Cys Leu Phe Phe Ile Phe Leu
                335                340                345

ttg tta ctt ctt tgc ctt att gtg gag aag gaa gaa cct gaa ccc aca 1347
Leu Leu Leu Leu Cys Leu Ile Val Glu Lys Glu Glu Pro Glu Pro Thr
                350                355                360

ttg agc ctc cat gtg gat ctg aac aaa gaa gtg tgt gct tat gaa gat 1395
Leu Ser Leu His Val Asp Leu Asn Lys Glu Val Cys Ala Tyr Glu Asp
365                370                375                380

acc ctc tgt taaaccacca atttcttgac atagagccag ccagcaggag 1444
Thr Leu Cys

tcatatataaa ctcaatttct cttaaaattt cgaatacatc ttcttgaaaa tccaaaaaaa 1504

aaaaaaaaa aaaaactcga g 1525

```

<210> 2
 <211> 383
 <212> PRT
 <213> Mus sp.

```

<400> 2
Met Ala Phe Val His Ile Arg Cys Leu Cys Phe Ile Leu Leu Cys Thr
  1                5                10                15

Ile Thr Gly Tyr Ser Leu Glu Ile Lys Val Asn Pro Pro Gln Asp Phe
                20                25                30

Glu Ile Leu Asp Pro Gly Leu Leu Gly Tyr Leu Tyr Leu Gln Trp Lys
                35                40                45

Pro Pro Val Val Ile Glu Lys Phe Lys Gly Cys Thr Leu Glu Tyr Glu
                50                55                60

Leu Lys Tyr Arg Asn Val Asp Ser Asp Ser Trp Lys Thr Ile Ile Thr
                65                70                75                80

Arg Asn Leu Ile Tyr Lys Asp Gly Phe Asp Leu Asn Lys Gly Ile Glu
                85                90                95

Gly Lys Ile Arg Thr His Leu Ser Glu His Cys Thr Asn Gly Ser Glu
                100                105                110

Val Gln Ser Pro Trp Ile Glu Ala Ser Tyr Gly Ile Ser Asp Glu Gly
                115                120                125

Ser Leu Glu Thr Lys Ile Gln Asp Met Lys Cys Ile Tyr Tyr Asn Trp
                130                135                140

Gln Tyr Leu Val Cys Ser Trp Lys Pro Gly Lys Thr Val Tyr Ser Asp

```

145 150 155 160
 Thr Asn Tyr Thr Met Phe Phe Trp Tyr Glu Gly Leu Asp His Ala Leu
 165 170 175
 Gln Cys Ala Asp Tyr Leu Gln His Asp Glu Lys Asn Val Gly Cys Lys
 180 185 190
 Leu Ser Asn Leu Asp Ser Ser Asp Tyr Lys Asp Phe Phe Ile Cys Val
 195 200 205
 Asn Gly Ser Ser Lys Leu Glu Pro Ile Arg Ser Ser Tyr Thr Val Phe
 210 215 220
 Gln Leu Gln Asn Ile Val Lys Pro Leu Pro Pro Glu Phe Leu His Ile
 225 230 235 240
 Ser Val Glu Asn Ser Ile Asp Ile Arg Met Lys Trp Ser Thr Pro Gly
 245 250 255
 Gly Pro Ile Pro Pro Arg Cys Tyr Thr Tyr Glu Ile Val Ile Arg Glu
 260 265 270
 Asp Asp Ile Ser Trp Glu Ser Ala Thr Asp Lys Asn Asp Met Lys Leu
 275 280 285
 Lys Arg Arg Ala Asn Glu Ser Glu Asp Leu Cys Phe Phe Val Arg Cys
 290 295 300
 Lys Val Asn Ile Tyr Cys Ala Asp Asp Gly Ile Trp Ser Glu Trp Ser
 305 310 315 320
 Glu Glu Glu Cys Trp Glu Gly Tyr Thr Gly Pro Asp Ser Lys Ile Ile
 325 330 335
 Phe Ile Val Pro Val Cys Leu Phe Phe Ile Phe Leu Leu Leu Leu Leu
 340 345 350
 Cys Leu Ile Val Glu Lys Glu Glu Pro Glu Pro Thr Leu Ser Leu His
 355 360 365
 Val Asp Leu Asn Lys Glu Val Cys Ala Tyr Glu Asp Thr Leu Cys
 370 375 380

<210> 3

<211> 1369

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (103)..(1245)

<400> 3

ggatccgcgc ggatgaaggc tatttgaagt cgccataacc tggtcagaag tgtgcctgtc 60

ggcggggaga gaggcaatat caagggtttta aatctcgag aa atg gct ttc gtt 114

Met Ala Phe Val

1

tgc ttg gct atc gga tgc tta tat acc ttt ctg ata agc aca aca ttt	162
Cys Leu Ala Ile Gly Cys Leu Tyr Thr Phe Leu Ile Ser Thr Thr Phe	
5 10 15 20	
ggc tgt act tca tct tca gac acc gag ata aaa gtt aac cct cct cag	210
Gly Cys Thr Ser Ser Ser Asp Thr Glu Ile Lys Val Asn Pro Pro Gln	
25 30 35	
gat ttt gag ata gtg gat ccc gga tac tta ggt tat ctc tat ttg caa	258
Asp Phe Glu Ile Val Asp Pro Gly Tyr Leu Gly Tyr Leu Tyr Leu Gln	
40 45 50	
tgg caa ccc cca ctg tct ctg gat cat ttt aag gaa tgc aca gtg gaa	306
Trp Gln Pro Pro Leu Ser Leu Asp His Phe Lys Glu Cys Thr Val Glu	
55 60 65	
tat gaa cta aaa tac cga aac att ggt agt gaa aca tgg aag acc atc	354
Tyr Glu Leu Lys Tyr Arg Asn Ile Gly Ser Glu Thr Trp Lys Thr Ile	
70 75 80	
att act aag aat cta cat tac aaa gat ggg ttt gat ctt aac aag ggc	402
Ile Thr Lys Asn Leu His Tyr Lys Asp Gly Phe Asp Leu Asn Lys Gly	
85 90 95 100	
att gaa gcg aag ata cac acg ctt tta cca tgg caa tgc aca aat gga	450
Ile Glu Ala Lys Ile His Thr Leu Leu Pro Trp Gln Cys Thr Asn Gly	
105 110 115	
tca gaa gtt caa agt tcc tgg gca gaa act act tat tgg ata tca cca	498
Ser Glu Val Gln Ser Ser Trp Ala Glu Thr Thr Tyr Trp Ile Ser Pro	
120 125 130	
caa gga att cca gaa act aaa gtt cag gat atg gat tgc gta tat tac	546
Gln Gly Ile Pro Glu Thr Lys Val Gln Asp Met Asp Cys Val Tyr Tyr	
135 140 145	
aat tgg caa tat tta ctc tgt tct tgg aaa cct ggc ata ggt gta ctt	594
Asn Trp Gln Tyr Leu Leu Cys Ser Trp Lys Pro Gly Ile Gly Val Leu	
150 155 160	
ctt gat acc aat tac aac ttg ttt tac tgg tat gag ggc ttg gat cat	642
Leu Asp Thr Asn Tyr Asn Leu Phe Tyr Trp Tyr Glu Gly Leu Asp His	
165 170 175 180	
gca tta cag tgt gtt gat tac atc aag gct gat gga caa aat ata gga	690
Ala Leu Gln Cys Val Asp Tyr Ile Lys Ala Asp Gly Gln Asn Ile Gly	
185 190 195	
tgc aga ttt ccc tat ttg gag gca tca gac tat aaa gat ttc tat att	738
Cys Arg Phe Pro Tyr Leu Glu Ala Ser Asp Tyr Lys Asp Phe Tyr Ile	
200 205 210	
tgt gtt aat gga tca tca gag aac aag cct atc aga tcc agt tat ttc	786
Cys Val Asn Gly Ser Ser Glu Asn Lys Pro Ile Arg Ser Ser Tyr Phe	
215 220 225	
act ttt cag ctt caa aat ata gtt aaa cct ttg ccg cca gtc tat ctt	834
Thr Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro Pro Val Tyr Leu	
230 235 240	

```

act ttt act cgg gag agt tca tgt gaa att aag ctg aaa tgg agc ata 882
Thr Phe Thr Arg Glu Ser Ser Cys Glu Ile Lys Leu Lys Trp Ser Ile
245                250                255                260

cct ttg gga cct att cca gca agg tgt ttt gat tat gaa att gag atc 930
Pro Leu Gly Pro Ile Pro Ala Arg Cys Phe Asp Tyr Glu Ile Glu Ile
                265                270                275

aga gaa gat gat act acc ttg gtg act gct aca gtt gaa aat gaa aca 978
Arg Glu Asp Asp Thr Thr Leu Val Thr Ala Thr Val Glu Asn Glu Thr
                280                285                290

tac acc ttg aaa aca aca aat gaa acc cga caa tta tgc ttt gta gta 1026
Tyr Thr Leu Lys Thr Thr Asn Glu Thr Arg Gln Leu Cys Phe Val Val
                295                300                305

aga agc aaa gtg aat att tat tgc tca gat gac gga att tgg agt gag 1074
Arg Ser Lys Val Asn Ile Tyr Cys Ser Asp Asp Gly Ile Trp Ser Glu
                310                315                320

tgg agt gat aaa caa tgc ttg gaa ggt gaa gac cta tcg aag aaa act 1122
Trp Ser Asp Lys Gln Cys Trp Glu Gly Glu Asp Leu Ser Lys Lys Thr
325                330                335                340

ttg cta cgt ttc ttg cta cca ttt ggt ttc atc tta ata tta gtt ata 1170
Leu Leu Arg Phe Trp Leu Pro Phe Gly Phe Ile Leu Ile Leu Val Ile
                345                350                355

ttt gta acc ggt ctg ctt ttg cgt aag cca aac acc tac cca aaa atg 1218
Phe Val Thr Gly Leu Leu Leu Arg Lys Pro Asn Thr Tyr Pro Lys Met
                360                365                370

att cca gaa ttt ttc tgt gat aca tga agactttcca tatcaagaga 1265
Ile Pro Glu Phe Phe Cys Asp Thr
                375                380

catgggtattg actcaacagt ttccagtcgat ggccaaatgt tcaatatgag tctcaataaa 1325

ctgaatttttt cttgcgaaaa aaaaaaaaaa aaatccgcgg atcc 1369

```

<210> 4

<211> 380

<212> PRT

<213> Homo sapiens

<400> 4

```

Met Ala Phe Val Cys Leu Ala Ile Gly Cys Leu Tyr Thr Phe Leu Ile
  1              5              10              15

Ser Thr Thr Phe Gly Cys Thr Ser Ser Ser Asp Thr Glu Ile Lys Val
                20              25              30

Asn Pro Pro Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr Leu Gly Tyr
                35              40              45

Leu Tyr Leu Gln Trp Gln Pro Pro Leu Ser Leu Asp His Phe Lys Glu
  50              55              60

```

Cys Thr Val Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly Ser Glu Thr
 65 70 75 80
 Trp Lys Thr Ile Ile Thr Lys Asn Leu His Tyr Lys Asp Gly Phe Asp
 85 90 95
 Leu Asn Lys Gly Ile Glu Ala Lys Ile His Thr Leu Leu Pro Trp Gln
 100 105 110
 Cys Thr Asn Gly Ser Glu Val Gln Ser Ser Trp Ala Glu Thr Thr Tyr
 115 120 125
 Trp Ile Ser Pro Gln Gly Ile Pro Glu Thr Lys Val Gln Asp Met Asp
 130 135 140
 Cys Val Tyr Tyr Asn Trp Gln Tyr Leu Leu Cys Ser Trp Lys Pro Gly
 145 150 155 160
 Ile Gly Val Leu Leu Asp Thr Asn Tyr Asn Leu Phe Tyr Trp Tyr Glu
 165 170 175
 Gly Leu Asp His Ala Leu Gln Cys Val Asp Tyr Ile Lys Ala Asp Gly
 180 185 190
 Gln Asn Ile Gly Cys Arg Phe Pro Tyr Leu Glu Ala Ser Asp Tyr Lys
 195 200 205
 Asp Phe Tyr Ile Cys Val Asn Gly Ser Ser Glu Asn Lys Pro Ile Arg
 210 215 220
 Ser Ser Tyr Phe Thr Phe Gln Leu Gln Asn Ile Val Lys Pro Leu Pro
 225 230 235 240
 Pro Val Tyr Leu Thr Phe Thr Arg Glu Ser Ser Cys Glu Ile Lys Leu
 245 250 255
 Lys Trp Ser Ile Pro Leu Gly Pro Ile Pro Ala Arg Cys Phe Asp Tyr
 260 265 270
 Glu Ile Glu Ile Arg Glu Asp Asp Thr Thr Leu Val Thr Ala Thr Val
 275 280 285
 Glu Asn Glu Thr Tyr Thr Leu Lys Thr Thr Asn Glu Thr Arg Gln Leu
 290 295 300
 Cys Phe Val Val Arg Ser Lys Val Asn Ile Tyr Cys Ser Asp Asp Gly
 305 310 315 320
 Ile Trp Ser Glu Trp Ser Asp Lys Gln Cys Trp Glu Gly Glu Asp Leu
 325 330 335
 Ser Lys Lys Thr Leu Leu Arg Phe Trp Leu Pro Phe Gly Phe Ile Leu
 340 345 350
 Ile Leu Val Ile Phe Val Thr Gly Leu Leu Leu Arg Lys Pro Asn Thr
 355 360 365
 Tyr Pro Lys Met Ile Pro Glu Phe Phe Cys Asp Thr
 370 375 380

<210> 5
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 5
ksrtccabk crctcca 17

<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 6
atagttaaac cattgccacc 20

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 7
ctccattcgc tccaaattcc 20

<210> 8
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 8
agtctatctt actttttactc g 21

<210> 9
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 9
catctgagca ataaatattc ac 22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/17103

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :A61K 38/16, 38/20

US CL :514/2, 12, 917; 424/145.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2, 12, 917; 424/145.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

West-US Patent full, STN via medline, embase, caplus, biosis. Search terms: interleukin 13 inhibit? or antagonis?, fibrosis, treatment or prevention

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CAPUT et al. Cloning and Characterization of a Specific Interleukin (IL)-13 Binding Protein Structurally Related to the IL-5 Receptor Alpha Chain. Journal Of Biological Chemistry. 12 July 1996. Vol.271, No.28, pages 16921-16926, see entire document.	1-30
A	HANCOCK et al. Production of Interleukin-13 by Alveolar Macrophages from Normal and Fibrotic Lung. American Journal of Respiratory Cell and Molecular Biology. January 1998. Vol.18, No.1, pages 60-67, especially see second and last paragraphs of page 65.	1-30



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

24 SEPTEMBER 2000

Date of mailing of the international search report

16 OCT 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

FOZIA HAMUD

Telephone No. (703) 308-0196